

J/ ψ Near-Threshold Photoproduction off the Proton and Neutron with CLAS12


RICHARD TYSON



University
of Glasgow

J/ψ Near Threshold Photoproduction

- ▶ CLAS12 operates close to the 8.2 GeV J/ψ ($c\bar{c}$ meson) photoproduction threshold.
- ▶ Near threshold, all the valence quarks of the nucleon are predicted to participate in the reaction compared to one or two at higher energies [2].
- ▶ [3] relates the nucleon gluonic form-factor to the t dependency of the differential cross section.
- ▶ CLAS12 will make a first measurement of J/ψ photoproduction on the neutron.



Measurements of the J/ψ total cross section as a function of the photon beam energy and theoretical predictions scaled to GlueX data [1].

- [1] A. Ali, et. al. (GlueX Collaboration), *Phys. Rev. Lett.* **123**, 072001 (2019).
[2] S. Brodsky, E. Chudakov, P. Hoyer, J. Laget, *Phys. Lett. B.* **498**, 23 (2001).
[3] L. Frankfurt, M. Strikman, *Phys. Rev. D.* **66**, 031502 (2002)



The CLAS12 Detector

- ▶ The CLAS12 Detector is located in Jefferson Lab's Hall B, in Newport News, Virginia.
- ▶ The GlueX detector is located in Hall D.



The CLAS12 Detector

- ▶ The CLAS12 Detector is located in Jefferson Lab's Hall B, in Newport News, Virginia.
- ▶ The recently upgraded CEBAF accelerator facility produces a 12 GeV electron beam, with beam energies up to 11 GeV delivered to Hall B.
- ▶ The Forward Detector has polar angle coverage of 5 to 35 degrees.
- ▶ The Central Detector has polar angle coverage of 35 to 125 degrees.



Experiment Overview


- ▶ J/ψ decays to a lepton pair, with l^+l^- denoting either e^+e^- or $\mu^+\mu^-$.
- ▶ CLAS12 took data with both a proton and a deuterium target offering several potential final states:

$$ep \rightarrow e' J/\psi p \rightarrow (e') l^+ l^- p$$

$$e p_{\text{bound}} \rightarrow e' J/\psi p \rightarrow (e') l^+ l^- p$$

$$e n_{\text{bound}} \rightarrow e' J/\psi n \rightarrow (e') l^+ l^- n$$


$$ed \rightarrow e' J/\psi d \rightarrow (e') l^+ l^- d$$



J/ψ quasi-real photoproduction
on a proton target



CLAS12 Forward Detector

- ▶ All final state particles are detected with the Forward Detector.
- ▶ The High Threshold Cherenkov Counter (HTCC) was built to identify electrons.
- ▶ The tracking system and Drift Chambers (DC) measure the charge and momentum of particles.
- ▶ The Forward Time Of Flight (FTOF) counters were designed to resolve charged hadrons.
- ▶ The Electromagnetic Calorimeters (PCAL and EC) are used to detect neutrals and identify electrons and muons.





Lepton Identification

- ▶ Electrons and positrons are required to produce a signal in the HTCC and high energy deposition in the calorimeter. Their main source of background is due to high momentum pions firing the HTCC.
- ▶ Muons are minimum ionising particles which we select with cuts on their energy deposition in the calorimeters. These are susceptible to a significant charged pion contamination.
- ▶ We refine leptons PID by training a machine learning classifier on variables from several CLAS12 detector subsystems to remove the pion background.
- ▶ The PID process is then reduced to a cut on the response of the classifier.




Hadron Identification

- ▶ For protons (and charged hadrons in general) a cut is made on the Beta versus Momentum parametrization.
- ▶ For neutrons we require a neutral charge. No further cuts were applied as there isn't any strong evidence of photon contamination.




Particle Corrections

- ▶ Radiative corrections for **electrons/positrons** add the momentum of **radiated photons**.
- ▶ **Neutrons** also produce **secondary clusters**. These are removed by taking the earliest neutral in a given sector.
- ▶ The reconstructed path length for neutrons is corrected for a more accurate calculation of the momentum.
- ▶ We apply fiducial cuts to remove e^+/e^- hits close to the edges of the PCAL where the shower is not fully contained within the calorimeter.
- ▶ Fiducial cuts in the drift chambers are applied to electrons, positrons, protons and muons by removing hits at the edge of the layers.





e- Sampling Fraction vs LV





Initial Event Selection

- ▶ To select only quasi-real photoproduction events regime we can minimize Q^2 .
- ▶ Similarly, we want the missing mass close to the mass of the scattered electron (which is effectively 0).
- ▶ The widths of these distributions can be parametrised as a function of the photon energy.
- ▶ The labels RG-A and RG-B refer to a proton or deuterium target respectively.





Di-Muon Invariant Mass

- ▶ Plotted here are the invariant mass distributions of:
 - ▶ $\mu^+\mu^-$ produced on a proton target.
 - ▶ $\mu^+\mu^-$ produced on a bound proton in the deuteron target.
- ▶ These are preliminary and produced with only a subset of all available data.





Di-Electron Invariant Mass

- ▶ Plotted here are the invariant mass distributions of:
 - ▶ e^+e^- produced on a bound neutron in the deuteron target.
 - ▶ e^+e^- produced on a bound proton in the deuteron target.
- ▶ The decrease in yield in the neutron channel compared to the proton channel is due to a lower detection efficiency for neutrons.
- ▶ These are preliminary and produced with only a subset of all available data.





ML Based Reaction ID

- ▶ We can extract a very clean J/ψ signal using a neural network trained on P/Theta/Phi of our final state particles.
- ▶ The training samples are:
 - ▶ J/ψ simulation.
 - ▶ Mismatched particles from different events of the above.
- ▶ In practice we need Bethe Heitler for normalisation.



Multiple Reactions

- ▶ Bethe Heitler has different kinematics to J/ψ and is removed by the classifier shown in the previous slide.
- ▶ We can train a classifier to ID 3 reactions:
 - ▶ J/ψ as before
 - ▶ Bethe Heitler photo/electroproduction
 - ▶ Mismatched particles from events in both above
- ▶ The principal can be applied to any analysis to disentangle different physical processes and sources of background.
- ▶ This is very much work in progress and we still have to validate this on data taken with CLAS12.



Conclusion



- ▶ The total and differential J/ψ photoproduction cross section provide unique insight on the J/ψ production mechanism and the nucleon gluonic form factor.
- ▶ CLAS12 will make a first measurement of the J/ψ photoproduction cross section ratio on proton and neutron.
- ▶ The analyses aiming for these measurements are ongoing and well developed.
- ▶ Next: total and differential cross sections for the proton and deuterium targets.



Backup Slides

P_c^+ resonances with CLAS12

- CLAS12 should be able to place upper limits on the branching fraction $B(P_c^+ \rightarrow J/\psi p)$ and $B(P_c^+ \rightarrow J/\psi n)$.






The $J/\psi p$ invariant mass distribution measured at the LHCb. Taken from:

R. Aaij, et. al. (LHCb Collaboration), *Phys. Rev. Lett.* **122**, 22 (2019).



P_c^+ Models

- ▶ Hadronic molecules: Weakly coupled charmed baryon and charmed meson.
- ▶ Hadro-charmonium states: compact bound $c\bar{c}$ state and light quarks.
- ▶ Quarks in a bag: Two tightly correlated di-quarks and an anti-quark.





Muon Identification

- ▶ Muons are minimum ionising particles which we select with cuts on their energy deposition in the CLAS12 calorimeters.
- ▶ Train a decision tree on MC data with training sample:
 - ▶ $\mu^+\mu^-$ past energy deposition cuts.
 - ▶ $\pi^+\pi^-$ past energy deposition cuts.





Muon ID Response

- ▶ The classifier output is given as a probability of being a signal event, called the response.
- ▶ The PID process is effectively reduced down to a cut on the response.
- ▶ This cut can be varied to study the systematic effect introduced by the classifiers.



Neutral ID


- ▶ We have neutrons with high beta, making it hard to cut on Beta to ID neutrals.
- ▶ From simulation however it doesn't look like we have large photon contamination past our exclusivity cuts.
- ▶ We therefore don't apply any neutral PID.




Neutron Path Corrections

- ▶ In the data reconstruction the neutron interaction point is set to the front face of the calorimeter when it should be within.
- ▶ Idea is to use $ep \rightarrow e' n \pi^+$ and compare calculated to reconstructed neutron path.
- ▶ Here we check three datasets:
 - ▶ RG-A spring2019 (10.2 GeV beam) – proton target
 - ▶ RG-B spring2019 (10.6 GeV beam) – deuterium target
 - ▶ RG-B spring2019 (10.2 GeV beam) – deuterium target

Missing Neutron Mass





ECin Reconstructed - Calculated Path




Neutron Path Corrections

- ▶ Correcting the path length means we can then correct the neutron Beta/momentum.
- ▶ We can use the same procedure to derive corrections for MC and data.





Secondary Neutrons

- ▶ Neutrons often produce secondary neutrons, due to:
 - ▶ real physical processes
 - ▶ Issues in reconstruction? Peak finding algorithm creates two clusters?
- ▶ This effect is seen in similar rates in both CLAS12 and MC (~30% of entries) with usual cuts and analysis procedures applied.



Secondary Neutrons

- ▶ Initial neutrons tend to occur first:
 - ▶ Earliest time when in same calorimeter layer
 - ▶ In layer closest to target when in different layers
- ▶ A couple of things left to iron out:
 - ▶ What happens when I have more than one particle in a sector (ie photons that are faster than neutrons)?
 - ▶ What happens if I have neutrons from another beam bunch?





Neutron Selection

- ▶ We have two selection procedures:
 - ▶ Only consider the **earliest neutral** in each sector as our main candidate.
 - ▶ Otherwise rank by χ^2 value calculated from exclusivity variables (MM^2/Q^2).


- ▶ χ^2 calculated as:

$$\chi^2 = \sum^{variables} \frac{(\text{observed} - \text{expected})^2}{\text{expected}}$$



$$ed \rightarrow (e')e^+e^-d$$

- ▶ Here the electron beam interacts with the deuteron as a whole.
- ▶ Deuteron ID isn't perfect at CLAS12 so we could possibly improve our statistics somewhat.
- ▶ This isn't a priority at the moment given low statistics.



ρ , ω and ϕ mesons

- ▶ Plotted here is the invariant mass of e^+e^- produced on a bound proton in the deuteron target.
- ▶ ϕ mesons are clearly resolved.
- ▶ ρ and ω mesons are unresolvable but clearly present.

