

Measurement of cross-section of the photoproduction of J/ψ on the proton

Pierre Chatagnon

13th of March 2024



Outline



Motivations



Analysis strategy and tools, Data and Monte-Carlo Samples



Results

J/ψ photoproduction at threshold

- The t -dependence of the cross-section allow to access gluon Gravitational Form Factors (GFFs), mass radius of the nucleon and gluon GPDs (under 2-gluon exchange assumption and no open-charm contributions)
- Model-dependent limit on the branching ration of the P_c pentaquark.

GlueX @ Jlab

Y-axis: $\sigma(\gamma p \rightarrow J/\psi p)$ [nb]

X-axis: \sqrt{s} [GeV]

Legend:

- GlueX (red circles with error bars)
- SLAC (blue squares with error bars)
- Cornell (green triangles with error bars)
- GPD (Ivanov, Sznajder, Szymanowski, and Wagner) (cyan shaded band)
- GPD and LQCD (Guo, Ji, and Liu) (orange dashed line)




Figure in Duran, B., Meiziani, Z.E., Joosten, S. et al. Determining the gluonic gravitational form factors of the proton. *Nature* 615, 813–816 (2023)

General analysis strategy

1) CLAS12 PID + Positron NN PID

$$ep \rightarrow (e')\gamma p \rightarrow (e')J/\psi p' \rightarrow (X)\overbrace{e^+e^-} p'$$

$$p_X = p_{beam} + p_p - p_{e^+} - p_{e^-} - p_{p'} \longrightarrow 2) |M_X^2| < 0.4 \text{ GeV}^2 \longrightarrow 3) Q^2 < 0.5 \text{ GeV}^2$$

Event selection

- Event topology:
 - exactly one electron in FD
 - exactly one positron in FD
 - exactly one proton
 - anything else
- HTCC and ECAL hits in the same sector
- HTCC lepton time within 2ns
- Lepton momenta > 1.7 GeV
- Proton in the FD
- Sampling Fraction > 0.15
- Lepton AI PID score > 0.05 (trained on pass 2 simulation)
- Exclusivity cuts:
 - $|M^2| < 0.4 \text{ GeV}^2$
 - $|Q^2| < 0.5 \text{ GeV}^2$


Data and MC samples

- Analysis on Pass 2 data. All *main* Fall 18 (Inbending and outbending) and Spring 19 runs are processed.
- Simulations are processed through OSG with pass 2 configuration
- The [QADB tool](#) is used to clean-up data and retrieve the accumulated charge per DST files
- The [RCDB interface of clas12root](#) is used to retrieve the beam current for each run
- Accumulated charge is computed per beam current for each configuration

Generator	Config / Beam currents / Charge					
	Fall 18 In.			Fall 18 Out.		Sp. 19
	45 nA 26.312 mC	50 nA 4.000 mC	55 nA 5.355 mC	40 nA 11.831 mC	50 nA 20.620 mC	50 nA 45.994 mC
Grape	8.2M each					6.7 M
TCSGen	2M each					1.5 M
JPsiGen	2M each					
JPsiGen (No rad.)	3M each					
Total of 24 MC samples and 3 Data samples						

Radiative corrections

- Inclusion of radiative effect is done in all generators according to formulas in: [Matthias Heller et al. Soft-photon corrections to the bethe-heitler process in the \$\gamma p \rightarrow l+l-p\$ reaction. PRD](#)
- The [JpsiGen](#), [TCSGen](#) generator with radiative effect are on Github, as well as an event converter for [Grape](#) ...not yet on OSG
- A full note on the algorithm is ready and will be included in the analysis note.
- The [work](#) was presented at the CLAS collaboration meeting in July 23.



Other analysis tools

https://github.com/PChatagnon/TCS_Analysis


Fiducial cuts/dead paddle cuts

- Pass I fiducial cuts on the PCAL (~ 8-9cm on V and W)
- Additional dead paddle cut, cross-check with Valerii Klimenko

Radiated photon correction

- Loop over photons in the event
- Add 4-vectors to the lepton if $\Delta\theta < 1.5$ deg.



Plots from Mariana Tenorio



- Not included in the following:
 - Energy loss / Energy corrections
 - Momentum smearing
 - Edge-based fiducial cuts

Results

Kinematic coverage and binning



Cross-section computation



Diagram illustrating the components of the cross-section σ_j :

$$\sigma_j = \mathcal{F}_j \cdot \mathcal{L} \cdot \omega_{cj} \cdot B_r \cdot \epsilon_j \cdot \epsilon_{Rad/j} \cdot N_{J/\psi j}$$

Annotations:

- $\mathcal{F}_j \cdot \mathcal{L}$: Number of photons and Number of targets (Red box)
- ω_{cj} : (Blue box)
- B_r : (Yellow box)
- ϵ_j : Reconstruction efficiency from MC (Dark blue box)
- $\epsilon_{Rad/j}$: Radiative corrections from MC (Orange box)
- $N_{J/\psi j}$: Branching ratio: 6% (Green box)

CLAS12 Preliminary





Number of J/Psi

$$\sigma_j = \frac{N_{J/\psi j}}{\mathcal{F}_j \cdot \mathcal{L} \cdot \omega_{cj} \cdot B_r \cdot \epsilon_j \cdot \epsilon_{Rad/j}}$$


- All data samples are combined and **fitted together**
- Gaussian + exponential background fit is used
- Systematic study is performed on the fit function

CLAS12 Preliminary



$$\sigma_j = \frac{N_{J/\psi j}}{\mathcal{F}_j \cdot \mathcal{L} \cdot \omega_{cj} \cdot B_r \cdot \epsilon_j \cdot \epsilon_{Rad/j}}$$

- $$\mathcal{L} = \frac{l \cdot \rho \cdot N_A \cdot C}{e}$$




Plots from Richard Tyson

Detection efficiency

$$\sigma_j = \frac{N_{J/\psi j}}{\mathcal{F}_j \cdot \mathcal{L} \cdot \omega_{cj} \cdot B_r \cdot \epsilon_j \cdot \epsilon_{Rad/j}}$$

- 1) From the data fit a second order polynomial background function is extracted
- 2) Events are generated according to this background function and added to the Jpsi signal MC sample
- 3) The obtained distribution is fitted with the same function as the data
- 4) The acceptance correction is then:

$$\epsilon_j = \frac{N_{J/\psi} |_{j/REC}}{N_{J/\psi} |_{j/RAD}}$$




Radiative correction


$$\sigma_j = \frac{N_{J/\psi j}}{\mathcal{F}_j \cdot \mathcal{L} \cdot \omega_{cj} \cdot B_r \cdot \epsilon_j \cdot \epsilon_{Rad/j}}$$

- 1) Jpsi samples without radiative effects are produced
- 2) The radiative correction is defined using the GEN kinematics as:

$$\epsilon_{Rad/j} = \frac{N_{J/\psi} \big|_{j/RAD}}{N_{J/\psi} \big|_{j/GEN}}$$






Preliminary cross-section as a function of E_γ




Selection cut systematics

- Every step of the analysis, except normalization factor, is repeated with different cuts:
 - Q^2 **DONE**
 - $|MM|^2$ **DONE**
 - Fit function **DONE**
 - Lepton momenta cut **To be done**
 - Lepton ID cut **To be done**
 - Proton PID **To be done**




Radiative correction effect

- The standard CS is extracted using the Radiated Jpsi MC samples and radiative correction
- The alternate is using non-radiated MC samples
- The effect is of the order of 10% (GlueX quoted 8.5%)




+ Closure test (Implemented but not presented here)

Bin volume correction



$$\left. \frac{d\sigma}{dt} \right|_j = \frac{N_{J/\psi/j}}{\mathcal{F}_j \cdot \mathcal{L} \cdot \omega_{c/j} \cdot B_r \cdot \epsilon_j \cdot \epsilon_{Rad/j} \cdot \boxed{\mathcal{V}_j \cdot \Delta t_j}}$$

- In practice is this readily done using integral of functions in root





t-dependence of the cross-section

- The t-dependent cross-section can be parametrized as:



$$\frac{d\sigma}{dt} = \frac{d\sigma}{dt} \Big|_0 \cdot \frac{1}{(1-t/m_s^2)^4}$$

- m_s can be interpreted as the mass radius of the proton. $\sqrt{\langle r_m^2 \rangle} = \frac{\sqrt{12}}{m_s}$
- Our results are not sensitive to the flattening at small E_γ and large t seen by GlueX.



Integrated t-dependent cross-section

- The integral of the t-dependent cross section is done bin-by-bin: $\sigma = \sum_j \left. \frac{d\sigma}{dt} \right|_j \cdot \Delta t_j$
- And compared to the total CS



- Good agreement between integrated t-dependent CS and E_γ -dependent CS

Take-aways and path going forward

I The J Ψ s analysis is at an advanced stage.


II Data and MC samples have been produced, the framework to analyze them is final.

III Some common tools remain to be developed and used in the analysis.

V **A release note will be ready by early April at the latest. An analysis note will be ready for the summer**

Back-up

Timeline for the tools and task for a dilepton publication



- On time for PID, Data processing and radiative corrections
- Still some tools required/preferred for the analysis (Momentum corrections/smearing)
- Still on track for analysis note submission by the summer

Data/MC normalization

- Each event is weighted by:

$$\omega = \frac{\mathcal{L} \cdot \sigma_{tot}}{nb_{GEN}} \quad \text{for generator providing integrated CS,}$$

$$\omega = \frac{\mathcal{L} \cdot w_{GEN}}{nb_{GEN}} \quad \text{for weighted generator.}$$

- Where the luminosity is obtained from target specification:



$$\mathcal{L} = \frac{l \cdot \rho \cdot N_A \cdot C \cdot Q}{e} = 1316.875 \cdot Q(\text{in mC})$$

Length of the target $l = 5 \text{ cm}$
Density of the target $\rho = 0.07 \text{ g/cm}^3$
Avogadro constant $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$
Unit charge $e = 1.6 \times 10^{-19} \text{ C}$
Conversion to pb $C = 10^{-36}$



https://clasweb.jlab.org/rungroups/tlc/wiki/images/e/e7/Normalization_MC_Data-5.pdf

Sampling fraction MC/Data mismatch

Inbending Fall 2018




Outbending Fall 2018




All Data fits


CLAS12 Preliminary




CLAS12 Preliminary



CLAS12 Preliminary




CLAS12 Preliminary






All Data fits

CLAS12 Preliminary




CLAS12 Preliminary




All Data fits

CLAS12 Preliminary



CLAS12 Preliminary




Lepton PID using AI


- Multiple evidences for large contamination from pions in the positron sample at high momenta ($P > 4.5$ GeV)
- We developed a PID algorithm to use on top of the EB PID for leptons (electron, positron, muon(sooon))
- Multivariate classifier using calorimeter responses only
 - Extension to Pass2 to the work that was done for the Pass1 TCS analysis
- One classifier per configuration and lepton flavor (6 in total)
- Soon available through Iguana
- Trained on simulation and validated on data

Work by Mariana Tenorio


Signal efficiency validation on data





Background rejection validation on data



Log scale results






Effect of using Frixione Flux



IV – Background modelisation and normalization


Comparison data/MC – Fall 2018 inbending

- Plotting conventions
 - Color-filled histograms are *stacked*, ie they show the total number of events with contributions for different channels “on top of each other”
 - Marker histograms are *not stacked* and simply superimposed









Overall strategy for background modelization

- 1) Event mixing
 - From data randomly select electron, positron, proton (from different events)
 - Construct kinematics and make sure they are within the region of interest ($M_{ee} > 2 \text{ GeV}$, $|MM|^2 < 0.4 \text{ GeV}^2$, $Q^2 < 2 \text{ GeV}^2$)
- 2) Reweight events to match data in the training region
- 3) Validate the weights on region B
- 4) Apply weights on region C and obtained BG-subtracted yields



Full comparison data/MC – Fall 2018 inbending (I)

$Q^2 \in [0.0, 0.5] \text{ GeV}^2$ Region C (Signal)



Training variables

Normalization factor

- Normalization factor can be computed as:

$$\omega_c = \frac{N_{Data} - N_{BG}}{N_{SIM\ BH}}$$

- Results:
 - Fall 2018 inbending – 68%
 - Spring 2019 inbending – 69%
 - Fall 2018 outbending – to be continued
- In the following, we use a normalization factor of 0.7

Spring 2019

CLAS12 Preliminary - Dilepton final state

