

Meson Spectroscopy at CLAS and CLAS12

An overview of selected results

Andrea Celentano
CLAS Collaboration

INFN-Genova




Introduction

- 1 Introduction
- 2 Meson spectroscopy at CLAS
- 3 The MesonEx experiment
- 4 Hidden-charm pentaquark search
- 5 Conclusions

Exotic mesons

QCD does not prohibit the existence of unconventional meson states such as hybrids ($q\bar{q}g$), tetraquarks ($q\bar{q}q\bar{q}$), and glueballs.



Exotic quantum numbers: $J^{PC} \neq q\bar{q}$


The discovery of states with manifest gluonic component, behind the CQM, would be the opportunity to directly “look” inside hadron dynamics.

Exotic quantum numbers would provide an **unambiguous** evidence of these states.

Lattice QCD calculations¹ provided a first hint on the spectrum and mass range of exotics.

Mass range: 1.4 GeV - 3.0 GeV

Lightest exotic is a 1^{-+} state.



¹J. J. Dudek et al, Phys. Rev. D82, 034508 (2010)

Exotic mesons photoproduction



Traditionally, meson spectroscopy was studied through different experimental techniques: **peripheral hadron production**, $N\bar{N}$ annihilation, ...

Photo-production measurements were limited by the lack of high-intensity, high-energy, high-quality photon beams.

Today, this limitation is no longer present.

Advantages:

- Photon spin: exotic quantum numbers are more likely produced by $S = 1$ probe
- Linear polarization: acts like a filter to disentangle the production mechanisms and suppress backgrounds
- Production rate: for exotics is expected to be comparable as for regular meson





Jefferson Laboratory - 6 GeV era

Jefferson Laboratory - today

Jefferson Laboratory (Newport News, VA, USA): home of the Continuous Electron Beam Accelerator Facility (CEBAF)

12-GeV e^- machine based on superconducting technology.

- 4 experimental Halls: A, B, C, D
- Multi-pass acceleration scheme, 2.2 GeV / pass
- Max. current: $\simeq 100\mu\text{A}$ / Hall (A and C)
- CW beam, $\simeq 100\%$ duty-cycle
- Beam polarization $\simeq 80\%$






The CLAS-g12 experiment

High-energy, high-statistics bremsstrahlung photon-beam experiment on LH_2 target

The g12 run period


- Summer 2008, $E_{e^-} = 5.715$ GeV, $I_{e^-} \simeq 60$ nA
- Tagged Bremsstrahlung photon-beam: 0.3-5.4 GeV, $L_{rad} = 10^{-4} X_0$
- 40-cm long LH_2 target
- Total number of recorded events $\simeq 26 \cdot 10^8$



CLAS detector

CLAS detector in Hall B at Jefferson Laboratory: almost 4π detector optimized for multi-particle final states


- Toroidal magnetic field (6 supercond. coils)
- Drift chambers (3 layers)
- Time-of-flight counters
- Electromagnetic calorimeters
 - $\sigma_E/E \simeq 10\%/ \sqrt{E}$
 - Angular coverage:
 $5^\circ \leq \theta \leq 45^\circ$
- Charged particle performances:
 - Acceptance: $8^\circ < \theta < 142^\circ$
 - Resolution: $\delta p/p \simeq 1\%$,
 $\delta\theta < 1 \text{ mrad}$



First measurement of the photoproduction reaction $\gamma p \rightarrow pa_2(1320)$

The $\gamma p \rightarrow p\pi^0\eta$ is a “golden channel” in meson spectroscopy: any P -wave resonance would unambiguously carry $J^{PC} = 1^{-+}$ exotic quantum numbers. A first measurement of this reaction in the multi-GeV energy range was recently completed using data from CLAS-g12.


- Both mesons were identified via their two-photon decay. A 4C kinematic fit was applied to the reaction $\gamma p \rightarrow p4\gamma$ events to ensure exclusivity.
- The *sPlot* technique was applied to isolate the reaction $\gamma p \rightarrow \pi^0\eta p$, using the invariant mass of the two photons from the η as control variable.
- The differential cross section $d^2\sigma/dtdM_{\pi^0\eta}$ was extracted in different E_γ and $-t$ kinematic bins



First measurement of the photoproduction reaction $\gamma p \rightarrow pa_2(1320)$

The $\gamma p \rightarrow p\pi^0\eta$ is a “golden channel” in meson spectroscopy: any P -wave resonance would unambiguously carry $J^{PC} = 1^{-+}$ exotic quantum numbers. A first measurement of this reaction in the multi-GeV energy range was recently completed using data from CLAS-g12.


- Both mesons were identified via their two-photons decay. A 4C kinematic fit was applied to the reaction $\gamma p \rightarrow p4\gamma$ events to ensure exclusivity.
- The $sPlot$ technique was applied to isolate the reaction $\gamma p \rightarrow \pi^0\eta p$, using the invariant mass of the two photons from the η as control variable.
- The differential cross section $d^2\sigma/dtdM_{\pi^0\eta}$ was extracted in different E_γ and $-t$ kinematic bins




First measurement of the photoproduction reaction $\gamma p \rightarrow pa_2(1320)$

The $\gamma p \rightarrow p\pi^0\eta$ is a “golden channel” in meson spectroscopy: any P -wave resonance would unambiguously carry $J^{PC} = 1^{-+}$ exotic quantum numbers. A first measurement of this reaction in the multi-GeV energy range was recently completed using data from CLAS-g12.

- Both mesons were identified via their two-photon decay. A 4C kinematic fit was applied to the reaction $\gamma p \rightarrow p4\gamma$ events to ensure exclusivity.
- The $sPlot$ technique was applied to isolate the reaction $\gamma p \rightarrow \pi^0\eta p$, using the invariant mass of the two photons from the η as control variable.
- The differential cross section $d^2\sigma/dtdM_{\pi^0\eta}$ was extracted in different E_γ and $-t$ kinematic bins



First measurement of the photoproduction reaction $\gamma p \rightarrow pa_2(1320)$




First measurement of the photoproduction reaction $\gamma p \rightarrow pa_2(1320)$

The limited statistics and acceptance prevented a full PWA of the final state. The dominant contribution of the $a_2(1320)$ to the total photo-production cross section was extracted in each E_γ and $-t$ kinematic bin through a fit to the $d^2\sigma/dtdM_{\pi^0\eta}$ observable via a resonance (a_2) + background model.

Most peculiar cross-section feature: dip at $-t \simeq 0.55 \text{ GeV}^2$ for both beam energies. From Regge phenomenology, considering the dominant ρ and ω exchanges (Mathieu, PRD 102, 2020):

$$A_{a_2} \propto (1 + \tau e^{i\pi\alpha(t)})\Gamma(1 - \alpha(t))$$


Our data also rule out other predictions for the $\gamma p \rightarrow pa_2$ photo-production cross section, based on other assumptions concerning the a_2 nature (for example, Xie et al. PRC 93, 2016)



$f_2(1270)$ photoproduction and decay via two π^0 channel

Exploiting the same $\gamma p \rightarrow p 4\gamma$ CLAS-g12 dataset, a high-statistics measurement of the $f_2(1270)$ photoproduction cross section on the proton was performed, exploiting the $f_2 \rightarrow \pi^0 \pi^0$ neutral decay channel. This acts as a "PWA-filter": no P-wave signals (i.e. no background from ρ production).


- In each E_{beam} and $-t$ beam, the f_2 yield was extracted from the $M_{\pi\pi}$ spectrum, performing a template fit to the f_2 signal and to the background (phase-space + f_0 tail)
- The cross section was determined from the measured f_2 yield, accounting for the CLAS acceptance/efficiency and for the luminosity.
- Results were compared with a prediction from a Regge-based model, finding a good agreement.



$f_2(1270)$ photoproduction and decay via two π^0 channel

Exploiting the same $\gamma p \rightarrow p 4\gamma$ CLAS-g12 dataset, a high-statistics measurement of the $f_2(1270)$ photoproduction cross section on the proton was performed, exploiting the $f_2 \rightarrow \pi^0 \pi^0$ neutral decay channel. This acts as a "PWA-filter": no P-wave signals (i.e. no background from ρ production).

- In each E_{beam} and $-t$ beam, the f_2 yield was extracted from the $M_{\pi\pi}$ spectrum, performing a template fit to the f_2 signal and to the background (phase-space + f_0 tail)
- The cross section was determined from the measured f_2 yield, accounting for the CLAS acceptance/efficiency and for the luminosity.
- Results were compared with a prediction from a Regge-based model, finding a good agreement.




Full PWA of $\gamma p \rightarrow n\pi^+\pi^+\pi^-$

Reaction $\gamma p \rightarrow \pi^+\pi^+\pi^-(n)$ - neutron identified via missing mass technique.
Focus on $E_\gamma > 4.4$ GeV to enhance meson resonances production.

C.Bookwalter (FSU)





- Clean 3π spectrum showing peaks of dominant resonances $a_2(1320)$ and $\pi_2(1670)$.
- Full PWA (17 waves): first time observation of $a_1(1260)$ in photoproduction.
- No signal of $\pi_1(1600)$ photoproduction.



Full PWA of $\gamma p \rightarrow n\pi^+\pi^+\pi^-$

Reaction $\gamma p \rightarrow \pi^+\pi^+\pi^-(n)$ - neutron identified via missing mass technique.
Focus on $E_\gamma > 4.4$ GeV to enhance meson resonances production.



- Clean 3π spectrum showing peaks of dominant resonances $a_2(1320)$ and $\pi_2(1670)$.
- Full PWA (17 waves): first time observation of $a_1(1260)$ in photoproduction.
- No signal of $\pi_1(1600)$ photoproduction.



Full PWA of $\gamma p \rightarrow n\pi^+\pi^+\pi^-$

Reaction $\gamma p \rightarrow \pi^+\pi^+\pi^-(n)$ - neutron identified via missing mass technique.
Focus on $E_\gamma > 4.4$ GeV to enhance meson resonances production.

- Clean 3π spectrum showing peaks of dominant resonances $a_2(1320)$ and $\pi_2(1670)$.
- Full PWA (17 waves): first time observation of $a_1(1260)$ in photoproduction.
- No signal of $\pi_1(1600)$ photoproduction.



From C. Bookwalter (FSU) PhD thesis. Full paper (A. Tsaris et al) submitted to PRL.

MesonEx (E12-005) in Hall-B at Jefferson Laboratory

Meson Spectroscopy program with quasi-real photons: low Q^2 electron scattering on a hydrogen target.




Goals:

- Measure the light-quarks mesons spectrum in the mass range 1.0 - 3.0 GeV
- Determine masses and properties of rare $q\bar{q}$ states
- Search for exotic mesons

Low Q^2 electron scattering:

- Provides a high-flux of high-energy, linearly polarized, quasi-real photons.
- Complementary and competitive to real photo-production
- Virtual photon kinematics and polarization determined event-by-event measuring scattered electron variables

Experimental technique: coincidence measurement between CLAS12 (final state hadrons) and Forward Tagger facility (low-angle scattered electron)



CLAS12 / Forward tagger detectors



CLAS12: multi-purpose, large acceptance, detector optimized for multi-particles final states (charged/neutrals)

- Nominal luminosity: $\mathcal{L} = 10^{35} \text{cm}^{-2} \text{s}^{-1}$
- Charged particles tracking: toroidal magnet + drift chambers system
- Particle ID: TOF, Cerenkov, RICH
- Neutral particles: lead/plastic scintillator calorimeter

Forward tagger: forward spectrometer optimized for detection of e^- scattered at low angle.


- **Lead-tungstate calorimeter (FT-Cal):** measure scattered electrons energy ($\sigma_E \simeq \%$)
- **Hodoscope (FT-Hodo):** distinguish photons from electrons.
- **Tracker (FT-Trck):** determine the electron scattering plane.

Nominal acceptance: $2.5^\circ < \theta_e < 4.5^\circ$,
 $0.5 < E_e(\text{GeV}) < 4.5$



MesonEx: expected results. Benchmark reaction: $\gamma p \rightarrow n \pi^+ \pi^+ \pi^-$ MC study










Isobar model for
3-pions production,
 $\sigma_{Tot} \simeq 10 \mu\text{barn}$



State	J^{PC}	L	Decay Mode
$a_1(1260)$	1^{++}	D	$\rho\pi$
$a_2(1320)$	2^{++}	D	$\rho\pi$
$\pi_2(1670)$	2^{-+}	P	$\rho\pi$
$\pi_2(1670)$	2^{-+}	F	$\rho\pi$
$\pi_2(1670)$	2^{-+}	S	$f_2\pi$
$\pi_2(1670)$	2^{-+}	D	$f_2\pi$
$\pi_1(1600)$	1^{-+}	P	$\rho\pi$

- 3π channel PWA feasible in MesonEx
- Sensitivity to $\pi_1(1600)$:
 $\sigma \geq 0.01 \sigma_{Tot}$
- Leakage contribution to exotic waves from others: $< 1\%$

Black: generated, Red: $t=-0.5 \text{ GeV}^2$, Blue: $t=-0.2 \text{ GeV}^2$ (D. Glazier, U. Glasgow)






MesonEx: expected results. Benchmark reaction $\gamma p \rightarrow p\pi^0\eta$ MC study

Ad-hoc model for reaction cross-section:

- Known resonances: $a_0(980)$, $a_2(1320)$, $a_2(1700)$
- Exotic contribution: $\pi_1(1400)$
- Large- $M_{\pi^0\eta}$: double-Regge exchange

Results:

- Non-exotic contributions properly reconstructed from PWA procedure
- Sensitivity to $\pi_1(1400)$ signal down to 5% of dominant $a_2(1320)$ signal






MesonEx: expected results. Benchmark reaction $\gamma p \rightarrow p\pi^0\eta$ MC study

Ad-hoc model for reaction cross-section:

- Known resonances: $a_0(980)$, $a_2(1320)$, $a_2(1700)$
- Exotic contribution: $\pi_1(1400)$
- Large- $M_{\pi^0\eta}$: double-Regge exchange

Results:



- Non-exotic contributions properly reconstructed from PWA procedure
- Sensitivity to $\pi_1(1400)$ signal down to 5% of dominant $a_2(1320)$ signal



MesonEx: π^0 photo-production

Motivation: day-0 analysis, involving only the FT detector. This will allow to solve the SLAC/GlueX tension on Σ .




- Reaction: $ep \rightarrow e'\pi^0(p)$: measure e' and two photons in FT, reconstruct proton via missing mass. $E_e = 10.6$ GeV
- Observables: Σ , $d\sigma/dt$ (also vs Q^2), σ_{TL} (not available in photoproduction).
- Status: analysis in progress exploiting the full CLAS12 RG-A 2018/2019 dataset (L. Biondo, Messina U.).



MesonEx: multi-particle final states

Day-1 analysis, core MesonEx program. All analysis are currently “work-in-progress”.

- 2π analysis (A. Thornton, Glasgow U.):
 $ep \rightarrow e' \pi^+ \pi^- (p)$
- 3π analysis (M. Nicol, York U.):
 $ep \rightarrow e' \pi^+ \pi^+ \pi^- (n)$
- $2K$ analysis (R. Wishart, Glasgow U.):
 $ep \rightarrow e' p K^+ K^- (p)$



LHCb hidden-charm pentaquark

LHCb in 2015 announced² the discovery of two exotic structures in the $J/\psi - p$ channel: $P_c(4380)$ and $P_c(4450)$, by measuring the decay $\Lambda_b^0 \rightarrow pJ/\psi K^-$.

They claimed that the minimum quark content is $c\bar{c}uud$.



Widths:

- $P_c(4450)$: $\Gamma = 39 \text{ MeV}$
- $P_c(4380)$: $\Gamma = 205 \text{ MeV}$

Quantum numbers (PWA most probable solution)

- $P_c(4450)$: $J_P = \frac{5}{2}^-$
- $P_c(4380)$: $J_P = \frac{3}{2}^+$

Although: "Acceptable solutions are also found for additional cases with opposite parity"



²Phys. Rev. Lett. **115**, 072001 (2015)

Hidden-charm pentaquark photo-production

A p - J/ψ resonance would appear as an s -channel resonance in the direct photo-production reaction:

$$\gamma p \rightarrow p J/\psi. \quad M_R = \sqrt{s} = M^2 + 2E_\gamma M$$

$$M_R \simeq 4.4 \text{ GeV} \rightarrow E_\gamma \simeq 10.1 \text{ GeV}$$

“Naive” cross-section estimate ingredients³:

- Breit-Wigner *elastic* cross-section
- Vector Meson Dominance

$$\sigma(W) = \frac{2J+1}{4} \frac{4\pi}{k_i^2} \frac{B_{in} B_{out} \Gamma^2/4}{(W-M_R)^2 + \Gamma^2/4}$$



Vector Meson Dominance:

$$B_{in} = (e/f_V)^2 B_{out} (k_{in}/k_{out})^{2L+1}$$

Cross-section estimate:

$$P_c(4380) : 1.5 \mu\text{barn} < \sigma_0/(B_{out}^2) < 50 \mu\text{barn}$$

$$P_c(4450) : 12 \mu\text{barn} < \sigma_0/(B_{out}^2) < 360 \mu\text{barn}$$



³ M. Karliner and J.L. Rosner, arXiv:1508.01496

Hall-B measurement



Use CLAS12 + Forward tagger detector for p - J/ψ quasi-real photo-production with two **complementary** techniques:

Untagged photo-production

- Scattered electron at $\theta_e \simeq 0^\circ$ not detected
- Measure final state p and e^+e^- from J/ψ decay with CLAS12
- Higher luminosity, lower W resolution.

Tagged photo-production

- Scattered electron detected in Forward Tagger, $2.5^\circ < \theta_e < 4.5^\circ$
- Measure in coincidence final state p and/or and e^+e^- from J/ψ decay with CLAS12
- p - J/ψ invariant mass W measured as missing mass on scattered e^- in Forward Tagger
- Lower luminosity, higher W resolution.






Hall-B measurement: untagged production

Reaction: $ep \rightarrow pe^+e^-(e')$. Results based on a limited data sample from 2018 run.

- Selection cuts based on transverse missing momentum, Q^2 , and missing mass. Optimal cut values determined using a boosted decision tree-based ML method.
- Clear evidence for light vector mesons production (ρ , ω , ϕ) - will be used as reference to check normalization and validate results.
- J/ψ signal is well visible.

J. Newton, Old Dominion University



Hall-B measurement: tagged production

Conclusions

- Experimental investigation of “exotic” hadrons is a powerful technique to answer to fundamental questions in QCD:
 - What is the origin of color confinement?
 - What is the role of gluons inside hadrons?
-
-
-



Backup slides

3 components:

- **Lead-tungstate calorimeter (FT-Cal):** measure the energy of scattered electrons with few % resolution.
- **Hodoscope (FT-Hodo):** distinguish photons from electrons.
- **Tracker (FT-Trck):** determine the electron scattering plane.

Nominal design parameters:

	Range
$E_{e'}$	0.5 - 4.5 GeV
$\theta_{e'}$	2.5° - 4.5°
$\phi_{e'}$	0° - 360°
E_γ	6.5 - 10.5 GeV
P_γ	70 - 10 %
Q^2	0.01 - 0.3 GeV^2 ($< Q^2 > 0.1 \text{ GeV}^2$)
W	3.6 - 4.5 GeV



The Forward Tagger Facility

3 components:

- **Lead-tungstate calorimeter (FT-Cal):** measure the energy of scattered electrons with few % resolution.
- **Hodoscope (FT-Hodo):** distinguish photons from electrons.
- **Tracker (FT-Trck):** determine the electron scattering plane.

Nominal design parameters:

	Range
$E_{e'}$	0.5 - 4.5 GeV
$\theta_{e'}$	2.5° - 4.5°
$\phi_{e'}$	0° - 360°
E_γ	6.5 - 10.5 GeV
P_γ	70 - 10 %
Q^2	0.01 - 0.3 GeV^2 ($< Q^2 > 0.1 \text{ GeV}^2$)
W	3.6 - 4.5 GeV

