

GANIL

**European Nuclear Physics
Conference 2025**

September 22-26

Caen, France

<https://indico.in2p3.fr/event/30430>

Measurement of beam-polarized
Deeply Virtual Compton Scattering
observables with ey detection @
CLAS12



CEBAF Large Acceptance Spectrometer



BY:

Juan Sebastian Alvarado

on behalf of the CLAS12 collaboration


EuNPC2025

23/09/2025



IJCLab - Orsay

université
PARIS-SACLAY



Jefferson Lab

C O N T E N T S

1

INTRODUCTION

Generalized Parton Distributions
Motivation

2

DATA SELECTION

BDT Classification
Background subtraction

3

DVCS MEASUREMENTS

Beam Spin Asymmetry
Cross-section

4

SUMMARY

1



INTRODUCTION

Generalized Parton Distributions


Motivation

1. INTRODUCTION



At low energy QCD, we describe the nucleon structure in terms of structure functions including:


1. Form Factors describe transverse position of partons
2. Parton Distribution Functions describe longitudinal momentum distributions
3. Generalized Parton Distributions (GPDs) correlates transverse position and longitudinal momentum distributions



GPDs encode information of the nucleon structure such as:

Nucleon Tomography

H.W. Lin, Phys. Rev. Lett. 127 (2021) 182001.



Contributions to the nucleon total spin.


X. Ji, Phy.Rev.Lett.78,610(1997)

$$\frac{1}{2} = \frac{1}{2} \Delta \Sigma + \Delta L + \Delta G$$

- Quark contribution is not the main contribution → Spin Crisis
- Quark's orbital angular momentum is accessed through GPDs
- Gluon contribution

Access to Gravitational Form Factors.

V. D. Burkert *et al.* Nature 557.7705 (2018): 396




- Mass/Energy distribution inside the nucleon
- Forces distribution.
- Nucleon mass radius
- Shear forces and pressure distribution

1. INTRODUCTION



The proton chiral-even GPDs can be accessed via electro-production of a real photon.



GPDs enters the DVCS-BH interference term through **Compton Form Factors**

$$\mathcal{F}(\xi, t) = \sum_q e_q^2 \left\{ \mathcal{P} \int_{-1}^1 dx \mathbf{F}^q(x, \xi, t) \left[\frac{1}{x - \xi} + \frac{1}{x + \xi} \right] - i\pi [\mathbf{F}^q(\xi, \xi, t) - \mathbf{F}^q(-\xi, \xi, t)] \right\}$$

which are measured in the Beam Spin Asymmetry

$$\text{BSA} = \frac{1}{P_{\text{beam}}} \frac{d^4\sigma^+ - d^4\sigma^-}{d^4\sigma^+ + d^4\sigma^-} \propto \Im \left\{ F_1 \mathcal{H} + \xi' (F_1 + F_2) \tilde{\mathcal{H}} - \frac{t}{4M_N^2} F_2 \mathcal{E} \right\},$$

For a spin $\frac{1}{2}$ particle, there are four chiral-even GPDs $F = H, E, \tilde{H}, \tilde{E}$

1. Detecting the proton is ideal, but by not detecting it:

- There is access to the small $-t=(p-p')^2$ region
- Larger statistics can be achieved, as the proton is not constrained by detector acceptance

2. However,

- There are background contributions from the whole SIDIS spectra.
- There are reduced options for cuts. Missing proton mass is the only exclusivity variable available

3. Therefore,

- We move to a ML approach for channel selection.
- It is first validated by including the proton information. Then applied to the no-proton case

2



DATA SELECTION



2. DATA SELECTION



Data taken by the **CLAS12** detector in fall-2018

- Polarized electron beam towards an unpolarized fixed proton target

CLAS12 is the large acceptance spectrometer located in the Hall B of the Thomas Jefferson National Accelerator Facility (Newport News, VA, USA)



KINEMATICS

- $W > 2 \text{ GeV}$
- $Q^2 > 1 \text{ GeV}^2$
- $q' > 2 \text{ GeV}$
- $e' > 1 \text{ GeV}$
- $(p' > 0.3 \text{ GeV})$

EXCLUSIVITY CUTS (if proton detected)

- $\Delta\phi = |\phi(p') - \phi(\gamma)| < 2^\circ$
- $\Delta t = |t(p') - t(\gamma)| < 2 \text{ GeV}^2$
- $P_{\text{miss}} < 1 \text{ GeV}$

$\phi(p')$ uses γ^* and p'

$\phi(\gamma)$ uses γ^* and γ

$$t(p') = (p - p')^2$$


$$t(\gamma) = \frac{Q^2 M_p + 2v M_p \left(v - \sqrt{v^2 + Q^2} \cos \theta_{\gamma^* \gamma} \right)}{\sqrt{v^2 + Q^2} \cos \theta_{\gamma^* \gamma} - v - M_p}$$

The event is built with the (e, γ , p) set with minimum missing $ep \rightarrow e\gamma p$ missing mass

2. DATA SELECTION

The main contamination channel is exclusive π^0 production

$$ep \rightarrow ep\pi^0 \rightarrow ep \gamma(\gamma)$$




Missing proton mass distribution

$$M_{e\gamma X}^2 = (e + p - e' - \gamma)^2$$


With proton

Without proton



In the case of no proton detection, there is contamination from **inclusive** π^0 production in SIDIS

2. CHANNEL SELECTION





Instead of straight cuts on exclusivity variables,
we optimize data selection with Boosted Decision Trees.

With proton

The following
variables are used
to train a BDT
due to its DVCS/DVMP
separation power.


W/out proton



After BDT
classification,
residual
background is
removed with
well known
methods


2. CHANNEL SELECTION

Data/MC comparison





Most of the background was removed, leading to a good agreement between simulation and data.

eyp/ey(p) comparison



When not requiring a proton:

- Increased statistics
- Access to smaller $-t$ values



The smaller t value, the better interpretation of DVCS in terms of GPDs!

3





DVCS MEASUREMENTS

Beam Spin Asymmetry

Cross-section

3. DVCS BSA



Christiaens, G., et al.
Phys. Rev. Lett. 130.21
(2023): 211902.

$$\mathbf{BSA} = \frac{1}{P_{\text{beam}}} \frac{d^4\sigma^+ - d^4\sigma^-}{d^4\sigma^+ + d^4\sigma^-}$$




$$\propto \text{Im} \left\{ F_1 \mathcal{H} + \xi' (F_1 + F_2) \tilde{\mathcal{H}} - \frac{t}{4M_N^2} F_2 \mathcal{E} \right\},$$

- Results compatible with previously published results.
- By not detecting the proton there is a better coverage in ϕ

3. DVCS BSA




BSA amplitude as a function of $-t$



There is sensitivity to $\cos\phi$ modulations at small $-t$

$$A_{LU}(\phi) \equiv \frac{p_1 \mathcal{P}_1(\phi) \mathcal{P}_2(\phi) \sin(\phi)}{1 + p_2 \cos(\phi) + p_3 \cos(2\phi)}$$





- The **ey(p)** selection gives access to the small $-t$ region
- Results compatible previous measurements

3. DVCS CROSS-SECTION

Juan Sebastian Alvarado, Université Paris-Saclay - IJCLab



Preliminary cross-section results
Case without proton information



Results with proton information will be published soon by other members of the collaboration



- GPDs are crucial for understanding the nucleon structure as they encode a variety of properties.
- DVCS is a golden channel for GPD studies due to its direct access to GPDs.
- The DVCS process was measured at Jefferson Lab, allowing the extraction of Beam Spin Asymmetry and cross-section measurements.
- While detecting the DVCS recoil proton is ideal, by not detecting it
 - There is a boost in statistics
 - There is access to a larger phase space
- Preliminary BSA and cross-section measurements were presented. Final results will be published soon.

THANKS
