

The proton's response to an external electromagnetic field challenges nuclear theory

R. Li¹, N. Sparveris^{1,*}, H. Atac¹, M.K. Jones², M. Paolone³, Z. Akbar⁴, V. Berdnikov⁵, D. Biswas⁶, M. Boer¹, A. Camsonne², J. -P. Chen², M. Diefenthaler², B. Duran¹, D. Dutta⁷, D. Gaskell², C. A. Gayoso⁸, O. Hansen², F. Hauenstein⁹, N. Heinrich¹⁰, W. Henry², T. Horn⁵, G. Huber¹⁰, S. Jia¹, S. Joosten¹¹, A. Karki⁷, S. Kay¹⁰, V. Kumar¹⁰, W. Li⁸, A. H. Liyanage⁶, D. Mack², S. Malace², P. Markowitz⁴, M. McCaughan², Z. E. Meziari¹¹, H. Mkrtchyan¹², C. Morean¹³, M. Muhoza⁵, A. Narayan¹⁴, B. Pasquini¹⁵, M. Rehfuss¹, B. Sawatzky², G. Smith², A. Smith¹⁶, R. Trotta⁵, C. Yero⁴, X. Zheng¹⁷, and J. Zhou¹⁶

List of affiliations*

ABSTRACT

Understanding how the visible matter in the universe arises from its elementary quark and gluon constituents is a central question for science. The visible world is founded on the proton, the only composite building block of matter that is stable in nature. Consequently, understanding the formation of matter relies on explaining the dynamics and the properties of the proton's bound state. A fundamental property of the proton involves the system's response to an external electromagnetic (EM) field. It is characterized by the EM polarizabilities¹ that describe how easily the charge and magnetization distributions inside the system are distorted by the EM field. When the polarizabilities are generalized to finite momentum transfer², by replacing the incoming real photon of the Compton scattering process with a space-like virtual photon, they map out the deformation of the quark densities in a proton subject to an EM field. They reveal vital information regarding the underlying system dynamics and provide a key for decoding the proton structure in terms of the theory of the strong interaction that binds quarks and gluons together. Here we report measurements of proton's electromagnetic generalized polarizabilities at low four-momentum transfer squared. We show evidence of an anomaly to the behaviour of proton's electric polarizability that contradicts the predictions of nuclear theory. The observed structure translates to a distinct signature in the spatial distribution of the induced polarization in the proton. The reported measurements suggest the presence of an unaccounted dynamical mechanism in the proton and present significant challenges to nuclear theory.

Explaining how the nucleons - protons and neutrons - emerge from the dynamics of their quark and gluon constituents is a central goal of modern nuclear physics. The importance of the question arises from the fact that the nucleons account for 99% of the visible matter in the universe. Moreover, the proton holds a unique role of being nature's only stable composite building block. The dynamics of quarks and gluons is governed by quantum chromodynamics (QCD), the theory of strong interaction. The application of perturbation methods renders QCD calculable at large energies and momenta - namely at high four-momentum transfer squared (Q^2) - and offers a reasonable understanding of the nucleon structure at that scale. Nevertheless, in order to explain the emergence of nucleon's fundamental properties from the in-

teractions of its constituents, the dynamics of the system have to be understood at long distances (or low Q^2), where the QCD coupling constant α_s becomes large and the application of perturbative QCD is not possible. The challenge arises from the fact that QCD is a highly nonlinear theory, since the gluons - the carriers of the strong force - couple directly to other gluons. Here, theoretical calculations can rely on lattice QCD³, a space-time discretization of the theory based on the fundamental quark and gluon degrees of freedom, starting from the original QCD Lagrangian. An alternative path is offered by effective field theories (EFTs), such as the chiral effective field theory⁴⁻⁶, which employ hadronic degrees of freedom and is based on the approximate and spontaneously broken chiral symmetry of QCD.

¹Temple University, Philadelphia, PA 19122, USA. ²Thomas Jefferson National Accelerator Facility, VA, USA. ³New Mexico State University, Las Cruces, NM 88003, USA. ⁴Florida State University, Tallahassee, FL 32306. ⁵Catholic University of America, Washington, DC 20064. ⁶Hampton University, Hampton, VA 23669. ⁷Mississippi State University, Miss. State, MS 39762. ⁸The College of William and Mary, Williamsburg, VA 23185. ⁹Old Dominion University, Norfolk, VA 23529. ¹⁰University of Regina, Regina, SK, Canada. ¹¹Argonne National Laboratory, Lemont, IL 60439. ¹²Artem Alikhanian National Laboratory, Yerevan, Armenia. ¹³University of Tennessee, Knoxville, TN 37996. ¹⁴Veer Kunwar Singh University, Arrah, Bihar 802301, India. ¹⁵University of Pavia, 27100 Pavia PV, Italy. ¹⁶Duke University, Durham, NC 27708. ¹⁷University of Virginia, Charlottesville, VA, 22904. **corresponding author: sparveri@temple.edu




Figure 1. Using virtual Compton scattering to measure the proton generalized polarizabilities

a) The experimental setup during the VCS (E12-15-001) experiment at Jefferson Lab. An electron beam impinges on a liquid hydrogen (red sphere) target. The interaction is mediated through a virtual photon (orange wavy line). The scattered electron and recoil proton are detected with two magnetic spectrometers, in coincidence. The real photon (green wavy line) that is produced in the reaction offers the electromagnetic perturbation and allows to measure the proton polarizabilities. **b)** The (undetected) real photon is identified through the reconstruction of the reaction's missing mass spectrum and allows the selection of the VCS events. **c)** The cross section of the VCS reaction measures the proton generalized polarizabilities. The dashed line denotes the Bethe Heitler+Born contributions to the cross section.

While steady progress has been made in recent years, we have yet to achieve a good understanding on how the nucleon properties emerge from the underlying dynamics of the strong interaction. In order to accomplish this, the theoretical calculations require experimental guidance and confrontation with precise measurements of the system's fundamental properties.

For a composite system, like the proton, the polarizabilities are fundamental structure constants, such as its size and shape. Listed among the system's primary properties at the Particle Data Group (PDG)¹, the two scalar polarizabilities - the electric, α_E , and the magnetic, β_M - can be interpreted as the response of the proton's structure to the application of an external electric or magnetic field, respectively. They describe how easily the charge and magnetization distributions inside the proton are distorted by the EM field and provide the net result on the system's spatial distributions. In order to measure the polarizabilities, one must generate an electric (\vec{E}) and a magnetic (\vec{H}) field. In the case of the proton, this is provided by the photons in the Compton scattering process. The two scalar polarizabilities appear as second order terms in the expansion of the real Compton Scattering (RCS)

amplitude in the energy of the photon

$$H_{eff}^{(2)} = -4\pi\left(\frac{1}{2}\alpha_E\vec{E}^2 + \frac{1}{2}\beta_M\vec{H}^2\right). \quad (1)$$

One can offer a simplistic description of the polarizabilities through the resulting effect of an electromagnetic perturbation applied to the nucleon constituents. An electric field moves positive and negative charges inside the proton in opposite directions. The induced electric dipole moment is proportional to the electric field, and the proportionality coefficient is the electric polarizability which quantifies the stiffness of the proton. On the other hand, a magnetic field acts in a different way on the quarks and on the pion cloud within the nucleon, giving rise to two different contributions in the magnetic polarizability, a paramagnetic and a diamagnetic, respectively. Compared to the atomic polarizabilities, which are of the size of the atomic volume, the proton electric polarizability α_E is much smaller than the volume scale of a nucleon¹. The small magnitude underlines the stiffness of the proton, as a direct consequence of the strong binding of the inner constituents, and indicates the intrinsic relativistic character of the system.

The generalization² of the two scalar polarizabilities in the four-momentum transfer space, $\alpha_E(Q^2)$ and $\beta_M(Q^2)$, is an extension of the static electric and magnetic polarizabilities obtained in RCS. They can be studied through measurements of the virtual Compton scattering (VCS) process² $\gamma^* p \rightarrow p\gamma$. The VCS is accessed experimentally through the $ep \rightarrow ep\gamma$ reaction. The definition of the reaction's kinematical parameters is given in the Methods section. Here, the incident real photon of the RCS process is replaced by a virtual photon. The virtuality of the incident photon (Q^2) sets the scale of the observation and allows one to map out the spatial distribution of the polarization densities in the proton, while the outgoing real photon provides the EM perturbation to the system. The meaning of the generalized polarizabilities (GPs) is analogous to that of the nucleon form factors. Their Fourier transform will map out the spatial distribution density of the polarization induced by an EM field. They probe the quark substructure of the nucleon and offer unique insight to the underlying nucleon dynamics. The interest on the GPs extends beyond the direct information that they provide on the dynamics of the system. They frequently enter as input parameters in various scientific problems. One such example involves the hadronic two-photon exchange corrections, which are needed for a precise extraction of the proton charge radius from muonic Hydrogen spectroscopy measurements⁷.




Figure 2. Feynman diagrams of the photon electroproduction

The mechanisms contributing to $ep \rightarrow ep\gamma$. The small circles represent the interaction vertex of a proton with a virtual photon and the ellipse the non-Born VCS amplitude.

In this work, we report on measurements of the VCS reaction at the Thomas Jefferson National Accelerator Facility (Jefferson Lab). The experiment accessed the region $Q^2 = 0.28 \text{ GeV}^2$ to 0.40 GeV^2 where the two scalar GPs are particularly sensitive to the nucleon dynamics. Of special interest is a puzzling measurement of $\alpha_E(Q^2)$ reported at $Q^2 = 0.33 \text{ GeV}^2$ ^{8,9} that involves a local enhancement of the electric GP, albeit of a large experimental uncertainty. This anomaly has been considered with reservation for many years. The theoretical calculations for the electric GP are

unable to account for such a feature and predict a monotonic fall-off with Q^2 . Recent experiments have not presented any supporting evidence of a local enhancement^{10,11}. In the reported work, we capitalize on the unique capabilities of the experimental setup at Jefferson Lab to conduct measurements of the scalar GPs with an unprecedented precision. The experiment exploits the sensitivity of the polarizabilities to the excited spectrum of the nucleon, that is e.g. different compared to the nucleon elastic form factors that describe only the ground state of the system. The measurements were conducted in the nucleon resonance region. This enables enhanced sensitivity to the polarizabilities compared to previous experiments that measured in the region of the pion production threshold. This has been previously exhibited e.g. in^{12,13}. Furthermore, cross section measurements were conducted for azimuthally symmetric kinematics in the photon angle, namely for $(\phi_{\gamma^* \gamma}, \pi - \phi_{\gamma^* \gamma})$. The measurement of the azimuthal asymmetry in the cross section enhances even further the sensitivity in the extraction of the polarizabilities, and suppresses part of the systematic uncertainties. Moreover, the $ep \rightarrow ep\pi^0$ reaction was measured, simultaneously with the $ep \rightarrow ep\gamma$ reaction. The pion electroproduction process is well understood in this kinematic regime, and its measurement offers a stringent, real-time normalization control to the measurement of the $ep \rightarrow ep\gamma$ cross section. Overall, a significant improvement was accomplished in the precision of the extracted generalized polarizabilities, compared to previous measurements.

The data were acquired in Hall C of Jefferson Lab during the VCS (E12-15-001) experiment. Electrons with energies of 4.56 GeV at a beam current up to $20 \mu\text{A}$ were produced by Jefferson Lab's Continuous Electron Beam Accelerator Facility (CEBAF) and were scattered from a 10 cm long liquid-hydrogen target. The Super High Momentum Spectrometer (SHMS) and the High Momentum Spectrometer (HMS) of Hall C were used to detect in coincidence the scattered electrons and recoil protons, respectively (see Fig. 1). Both spectrometers are equipped with similar detector packages, including a set of scintillator planes, that were used to form the trigger and to provide time-of-flight information and a pair of drift chambers used for tracking. The coincidence time was determined as the difference in the time-of-flight between the two spectrometers, accounting for path-length variation corrections from the central trajectory and for the individual start-times. The experimental setup offered a $\sim 1 \text{ ns}$ (FWHM) resolution in the coincidence timing spectrum. The random coincidences were subtracted using the side (accidental) bands of the coincidence time spectrum. The events of the exclusive reaction $ep \rightarrow ep\gamma$ (see Fig. 2) were identified from the missing-mass reconstruction, through a selection cut around the photon peak in the missing-mass-squared spectrum. Data were taken with an empty target in order to account for the background contributions from the target walls. Elastic scattering measurements with a proton target were performed throughout the experiment for




Figure 3. Cross section measurements of the VCS reaction

a) Cross section measurements for in-plane kinematics at $Q^2 = 0.28 \text{ GeV}^2$. Results are shown for different bins in the total c.m. energy of the (γp) system, W . **b)** Measurements for in-plane kinematics at $Q^2 = 0.33 \text{ GeV}^2$. **c)** Measurements for in-plane kinematics at $Q^2 = 0.40 \text{ GeV}^2$. Top and bottom panels correspond to $\phi_{\gamma^* \gamma} = 180^\circ$ and $\phi_{\gamma^* \gamma} = 0$, respectively. The solid curve shows the Dispersion Relations (DR) fit for the two scalar generalized polarizabilities. The dashed curve shows the Bethe-Heitler plus Born-VCS (BH+Born) cross section. The error bars correspond to the total uncertainty, at the 1σ or 68% confidence level.

calibration and normalization studies. The measurement of the absolute VCS cross section, $\sigma \equiv d^5\sigma/dE'_e d\Omega'_e d\Omega_{cm}$, requires the determination of the five-fold solid angle, where dE'_e , $d\Omega'_e$ is the differential energy and solid angle of the scattered electron in the laboratory frame and $d\Omega_{cm}$ is the differential solid angle of the proton in the center-of-mass. The experimental acceptance is calculated through the Hall C Monte Carlo simulation program, SIMC, which integrates the beam configuration, target geometry, spectrometer acceptances, resolution effects, energy losses and radiative corrections. The cross section results for in-plane kinematics are presented in Fig. 3. The measurements are shown for different bins in the total c.m. energy of the (γp) system, W . They span across an extended range of $\theta_{\gamma^* \gamma}$ and avoid the kinematics dominated by the Bethe-Heitler process where the polarizability effect is suppressed. The complete dataset of the measured cross sections is included in the Extended

Data tables.

The cross section of the $ep \rightarrow ep\gamma$ process observes the photon that is emitted by either the lepton, known as the Bethe-Heitler (BH) process, or by the proton, the fully virtual Compton scattering (FVCS) process, as shown in Fig. 2. The FVCS amplitude can in-turn be decomposed into a Born contribution, with the intermediate state being the nucleon, and a non-Born contribution that carries the physics of interest and is parametrized by the GPs. The BH and the Born-VCS contributions are well known, calculable in terms of the proton electromagnetic form factors that are precisely measured from elastic electron scattering. We extract the GPs from the measured cross sections through a fit that employs the dispersion relations (DR) model¹⁴⁻¹⁶ for VCS. In the DR formalism, the two scalar GPs enter unconstrained and can be adjusted as free parameters, while the proton electromagnetic form factors are introduced as an input. The




Figure 4. The generalized polarizabilities of the proton

a) The electric generalized polarizability measured in this experiment (red symbols). The world data^{8–13,31–34} are shown with open-symbols. Circle and box symbols indicate results following the Dispersion-Relations and Low-Energy-Expansion analysis, respectively. The theoretical calculations of BChPT¹⁷, NRQCM²⁸, LSM²³, ELM²⁵ and DR^{14–16} are also shown. **b)** The magnetic generalized polarizability. The definition of symbols and curves are the same as in (a). **c)** Induced polarization in the proton when submitted to an EM field as a function of the transverse position with photon polarization along the x axis. **d)** The proton electric polarizability radius $r_{\alpha_E} \equiv \sqrt{\langle r_{\alpha_E}^2 \rangle}$ derived from this work (red point). The measurements of the proton charge radius r_E ^{35–42} (blue points) are shown for comparison. The error bars correspond to the total uncertainty, at the 1σ or 68% confidence level.

experimental cross sections are compared to the DR model predictions for all possible values for the two GPs, and the $\alpha_E(Q^2)$ and $\beta_M(Q^2)$ are fitted by a χ^2 minimization. The extracted electric and magnetic GPs are shown in Fig. 4. We observe a local enhancement of $\alpha_E(Q^2)$ in the measured region. For $\beta_M(Q^2)$, we find a smooth Q^2 -dependence and the near-cancellation of the paramagnetic and the diamagnetic contributions in the proton at $\sim Q^2 = 0.4 \text{ GeV}^2$. From the theory standpoint, the GPs have been calculated employing a variety of approaches, namely chiral effective field theories^{17–22}, the linear σ -model^{23,24}, the Effective Lagrangian Model²⁵, relativistic²⁶ and nonrelativistic^{27,28} constituent quark models. Lattice-QCD calculations are for the moment limited to polarizabilities in RCS²⁹. The current theoretical understanding suggests that $\alpha_E(Q^2)$ decreases monotonically

as the distance scale becomes smaller, namely with increasing Q^2 . The $\beta_M(Q^2)$ is expected to have a smaller magnitude, relative to $\alpha_E(Q^2)$. This can be explained by the competing paramagnetic and diamagnetic contributions in the proton, which largely cancel. In some theoretical calculations, $\beta_M(Q^2)$ is predicted to go through a maximum before decreasing. This last feature is typically explained by the dominance of diamagnetism due to the pion cloud at long distance (or small Q^2) and the dominance of paramagnetism due to a quark core at short distance scales. The theoretical predictions for the two generalized polarizabilities vary noticeably in magnitude. The reported measurements impose strict constraints and provide new input to the theory. The remarkable observation involves the measured $\alpha_E(Q^2)$ anomaly. It manifests as a distinct local deviation from the

single-dipole behavior that describes the rest of the world data, as discussed in^{10,11}. It contradicts all the theoretical calculations, that unanimously predict a smooth fall-off as a function of Q^2 . The theory is missing a dynamical element that can explain how a local enhancement of the system's electric polarizability can emerge as the distance scale becomes smaller, namely where the quark degrees of freedom acquire an increasingly prominent role in the dynamics of the system. The data suggest the presence of a dynamical mechanism in the proton that is currently not accounted for in the theory.

From the measurements of the generalized polarizabilities, we derive the spatial deformation of the quark distributions in the proton subject to the influence of an external electromagnetic field³⁰ (see Methods for details). This follows effectively an extension of the formalism to extract the light-front quark charge densities⁴³ from the proton form factor data. First, we derive an accurate Q^2 -parametrization of the polarizabilities from a fit to the experimental data. From that, we extract the induced polarization in the proton following³⁰. As shown in Fig. 4(c), we observe that the enhancement of $\alpha_E(Q^2)$ is translated to a distinct structure in the spatial distribution of the induced polarization in the proton. The distribution follows a change of sign around ~ 0.25 fm and exhibits a secondary maximum in the amplitude around ~ 0.35 fm. A primary measure that quantifies the extension of a spatial distribution is the mean square radius. The mean square electric polarizability radius of the proton $\langle r_{\alpha_E}^2 \rangle$ is related to the slope of the electric GP at $Q^2 = 0$

$$\langle r_{\alpha_E}^2 \rangle = \frac{-6}{\alpha_E(0)} \cdot \left. \frac{d}{dQ^2} \alpha_E(Q^2) \right|_{Q^2=0}. \quad (2)$$

We determine the slope of $\alpha_E(Q^2)$ at $Q^2 = 0$ from fits to the world-data, using a group of functional forms that can provide a reliable fit (see Methods for details). For $\alpha_E(0)$, we adopt the most recent measurement from³¹. For the mean square electric polarizability radius we find $\langle r_{\alpha_E}^2 \rangle = 1.28 \pm 0.24$ fm². This value is considerably larger compared to the mean square charge radius of the proton, $\langle r_E^2 \rangle \sim 0.7$ fm²¹ (see Fig. 4(d)). The dominant contribution to this effect is expected to arise from the deformation of the mesonic cloud in the proton under the influence of an external EM field. We derive the mean square magnetic polarizability radius from the magnetic polarizability measurements, following a procedure that is equivalent to the extraction of the mean square electric polarizability radius (see Methods for details) and we find that $\langle r_{\beta_M}^2 \rangle = 0.63 \pm 0.31$ fm².

In conclusion, we have studied proton's response to an external electromagnetic field and its dependence on the distance scale within the system. We show evidence of a local enhancement in proton's electric generalized polarizability that the nuclear theory cannot explain. The reported data suggest the presence of a dynamical mechanism in the system that is currently not accounted for in the theory. Proton

has the unique role of being nature's only stable composite building block. Consequently, the observed anomaly in the system's response to an electromagnetic field comes with a unique scientific interest. It calls for further measurements so that the underlying dynamics can be mapped with precision and highlights the need for an improved theory so that a fundamental property of the proton can be reliably described.

References

1. Zyla, P. A. *et al.* Review of Particle Physics. *Prog. Theor. Exp. Phys.* **2020**, 083C01 (2020).
2. Fonvieille, H., Pasquini, B. and Sparveris, N. Virtual Compton Scattering and Nucleon Generalized Polarizabilities. *Prog. Part. Nucl. Phys.* **113**, 103754 (2020).
3. Wilson, K. G. Confinement of quarks *Phys. Rev. D* **10**, 2445 (1974).
4. Weinberg, S. Nuclear forces from chiral lagrangians *Phys. Lett. B* **251**, 288 (1990).
5. Weinberg, S. Effective chiral lagrangians for nucleon-pion interactions and nuclear forces *Nucl. Phys. B* **363**, 3 (1991).
6. Weinberg, S. Three-body interactions among nucleons and pions *Phys. Lett. B* **295**, 114 (1992).
7. Pasquini, B. and Vanderhaeghen, M. Dispersion Theory in Electromagnetic Interactions *Annu. Rev. Nucl. Part. Sci.* **68**, 75-103 (2018).
8. Roche, J. *et al.* First Determination of Generalized Polarizabilities of the Proton by a Virtual Compton Scattering Experiment *Phys. Rev. Lett.* **85**, 708-711 (2000).
9. Janssens, P. *et al.* A new measurement of the structure functions PLL-PTT/ ϵ and PLT in virtual Compton scattering at $Q^2 = 0.33(\text{GeV}/c)^2$ *Eur. Phys. J.* **A37**, 1-8 (2008)
10. Beričič, J. *et al.* New Insight in the Q^2 Dependence of Proton Generalized Polarizabilities *Phys. Rev. Lett.* **123**, 192302 (2019)
11. Fonvieille, H. *et al.* Measurement of the generalized polarizabilities of the proton at intermediate Q^2 *Phys. Rev. C* **103**, 025205 (2021)
12. Laveissiere, G. *et al.* Measurement of the Generalized Polarizabilities of the Proton in Virtual Compton Scattering at $Q^2 = 0.92$ and 1.76 GeV² *Phys. Rev. Lett.* **93**, 122001 (2004)
13. Fonvieille, H. *et al.* Virtual Compton scattering and the generalized polarizabilities of the proton at $Q^2 = 0.92$ and 1.76 GeV² *Phys. Rev. C* **86**, 015210 (2012)
14. Pasquini, B., Drechsel, D., Gorchtein, M., Metz, A., and Vanderhaeghen, M. Dispersion relation formalism for virtual Compton scattering and the generalized polarizabilities of the nucleon *Phys. Rev. C* **62**, 052201(R) (2000).

15. Pasquini, B., Gorchtein, M., Drechsel, D., Metz, A. and Vanderhaeghen, M. Dispersion relation formalism for virtual Compton scattering off the proton *Eur. Phys. J. A* **11**, 185 (2001).
16. Drechsel, D., Pasquini, B. and Vanderhaeghen, M. Dispersion relations in real and virtual Compton scattering *Phys. Rep.* **378**, 99 (2003).
17. Lensky, V., Pascalutsa, V. and Vanderhaeghen, M. Generalized polarizabilities of the nucleon in baryon chiral perturbation theory *Eur. Phys. J. C* **77**, 119 (2017).
18. Hemmert, T. R., Holstein, B., Knochlein, G., Scherer, S. Virtual Compton scattering off the nucleon in chiral perturbation theory *Phys. Rev. D* **55**, 2630 (1997).
19. Hemmert, T. R., Holstein, B., Knochlein, G., Scherer, S. Generalized polarizabilities and the chiral structure of the nucleon *Phys. Rev. Lett.* **79**, 22 (1997).
20. Hemmert, T. R., Holstein, B., Knochlein, G., Drechsel, D. Generalized polarizabilities of the nucleon in chiral effective theories *Phys. Rev. D* **62**, 014013 (2000).
21. Kao, C. W., Vanderhaeghen, M. Generalized spin polarizabilities of the nucleon in heavy baryon chiral perturbation theory at next-to-leading order *Phys. Rev. Lett.* **89**, 272002 (2002).
22. Kao, C. W., Pasquini, B., Vanderhaeghen, M. New predictions for generalized spin polarizabilities from heavy baryon chiral perturbation theory *Phys. Rev. D* **92**, 119906 (2015).
23. Metz, A., Drechsel, D. Generalized polarizabilities of the nucleon studied in the linear sigma model *Z. Phys. A* **356**, 351 (1996).
24. Metz, A., Drechsel, D. Generalized polarizabilities of the nucleon studied in the linear sigma model. 2 *Z. Phys. A* **359**, 165 (1997).
25. Korchin, A. and Scholten, O. Nucleon polarizabilities in virtual Compton scattering *Phys. Rev. C* **58**, 1098 (1998)
26. Pasquini, B. and Salme, G. Nucleon generalized polarizabilities within a relativistic constituent quark model *Phys. Rev. C* **57**, 2589 (1998).
27. Liu, G. Q., Thomas, A. W., Guichon, P. A. M. Virtual Compton scattering from the proton and the properties of nucleon excited states *Aust. J. Phys.* **49**, 905 (1996).
28. Pasquini, B., Scherer, S. and Drechsel, D. Generalized polarizabilities of the proton in a constituent quark model revisited *Phys. Rev. C* **63**, 025205 (2001).
29. Detmold, W., Tiburzi, B., Walker-Loud, A. Extracting Nucleon Magnetic Moments and Electric Polarizabilities from Lattice QCD in Background Electric Fields *Phys. Rev. D* **81**, 054502 (2010).
30. Gorchtein, M., Lorce, C., Pasquini, B., Vanderhaeghen, M. Light-front interpretation of Proton Generalized Polarizabilities *Phys. Rev. Lett.* **104**, 112001 (2010).
31. Mornacchi, E. *et al.* Measurement of Compton scattering at MAMI for the extraction of the electric and magnetic polarizabilities of the proton *arXiv:2110.15691*, (2010)
32. Bourgeois, P. *et al.* Measurements of the generalized electric and magnetic polarizabilities of the proton at low Q^2 using the VCS reaction *Phys. Rev. Lett.* **97**, 212001 (2006)
33. Bourgeois, P. *et al.* Measurements of the generalized electric and magnetic polarizabilities of the proton at low Q^2 using the virtual Compton scattering reaction *Phys. Rev. C* **84**, 035206 (2011)
34. Blomberg, A. *et al.* Virtual Compton Scattering measurements in the nucleon resonance region. *Eur. Phys. J. A* **55**, 182 (2019).
35. Pohl, R. *et al.* The size of the proton *Nature* **466**, 213-216 (2010)
36. Antognini, A. *et al.* Proton structure from the measurement of 2S–2P transition frequencies of muonic hydrogen *Science* **339**, 417–420 (2013)
37. Xiong, W. *et al.* A small proton charge radius from an electron–proton scattering experiment *Nature* **575**, 147–150 (2019)
38. Fleurbaey, H. *et al.* New measurement of the 1S–3S transition frequency of hydrogen: contribution to the proton charge radius puzzle *Phys. Rev. Lett.* **120**, 182001 (2018)
39. Bezginov, N. *et al.* A measurement of the atomic hydrogen Lamb shift and the proton charge radius *Science* **365**, 1007 (2019)
40. Bernauer, J. C. *et al.* High-Precision Determination of the Electric and Magnetic Form Factors of the Proton *Phys. Rev. Lett.* **105**, 242001 (2010)
41. Atac, H. *et al.* Charge radii of the nucleon from its flavor dependent Dirac form factors *Eur. Phys. J. A* **57**, 65 (2021)
42. Grinin, A. *et al.* Two-photon frequency comb spectroscopy of atomic hydrogen *Science* **370**, 1061–1066 (2020)
43. Atac, H. *et al.* Measurement of the neutron charge radius and the role of its constituents *Nat. Commun.* **12**, 1759 (2021)

Methods

Experimental setup and measurements. For the measurement of the $ep \rightarrow ep\gamma$ reaction, electrons with energies of 4.56 GeV at a beam current up to $20 \mu\text{A}$ were produced by Jefferson Lab's Continuous Electron Beam Accelerator Facility (CEBAF). The electrons were scattered from a 10 cm long liquid-hydrogen target at a temperature of 19 K. The thickness of the aluminum target cell at the entrance and exit is 0.130 (12) mm and 0.188 (13) mm, respectively. For every kinematical setting, data were taken with a target made of two aluminum foils located at the positions of the cryotarget entrance and exit windows, each having a thickness of 0.6463 (8) mm, in order to subtract the background contributions emerging from the target walls by scaling the thicknesses of the two targets. The scattered electron and recoil proton of the reaction are detected with two magnetic spectrometers, in coincidence. The outgoing photon of the VCS process is identified through the reconstructed missing mass spectrum. The polar angle $\theta_{\gamma^* \gamma}$ of the VCS reaction is defined as the center-of-mass (c.m.) polar angle of the real photon with respect to the momentum transfer direction. The azimuthal angle of the reaction $\phi_{\gamma^* \gamma}$ defines the angle between the plane of the two (incoming and scattered) electrons and the photon-proton plane. The four-momentum of the outgoing photon, denoted by \mathbf{q}' , is reconstructed as $\mathbf{q}' = \mathbf{k} + \mathbf{p} - \mathbf{k}' - \mathbf{p}'$, where \mathbf{k} and \mathbf{p} are the four-momenta of the incoming electron and the target proton, while \mathbf{k}' and \mathbf{p}' are the four-momenta of the final electron and proton, respectively. The four-momentum of the virtual photon is $\mathbf{q} = \mathbf{k} - \mathbf{k}'$, with $Q^2 \equiv -\mathbf{q}^2$.

The beam properties were monitored throughout the experiment with the Hall C beam diagnostic elements. The beam position monitors (BPMs), that consist of a 4-wire antenna array of open ended thin wire striplines tuned to the RF frequency of the beam, were used to determine the position and the direction of the beam on the experimental target point. The beam current monitors (BCMs), a set of resonant-cavity based beam-current monitors and a parametric current transformer monitor, were used for the continuous non-intercepting beam current measurements. The beam size was measured by using harp scanners, which moved a thin wire through the beam. The beam was spread over a $2 \times 2 \text{ mm}^2$ area to avoid overheating the target. The beam energy was determined with an uncertainty of 0.06% by measuring the bend angle of the beam, on its way into Hall C, as it traversed the Hall C arc dipole magnets. The total accumulated beam charge was determined with 0.5% uncertainty. The liquid-hydrogen target density is determined within 0.7% and the target length uncertainty is 0.26%.

Two magnetic spectrometers, the Super High Momentum Spectrometer (SHMS) and the High Momentum Spectrometer (HMS) were used to detect in coincidence the scattered electrons and recoil protons, respectively. Both spectrometers involve a series of superconducting magnets, including quadrupoles and dipoles, followed by a set of particle detectors. The dipole magnets deflect charged particles vertically

as they enter the detector huts, while the quadrupole magnets optimize the flux of the charged particles entering the dipole magnet and focus the orbits of the charged particles into the detector huts. The two spectrometers are equipped with similar detector packages, with some differentiation due to the different momentum ranges of the spectrometers. The SHMS is also equipped with a Pb-glass calorimeter⁴⁴ that can serve as a particle identification detector. A pair of drift chambers, each with 6 wire planes separated by about a meter, was used to provide the tracking of the detected particles. The uncertainty in the determination of the tracking efficiency was 0.5% and 1% for the SHMS and the HMS, respectively. A set of hodoscope planes was used to form the trigger and to provide time-of-flight information. The time-of-flight in the HMS spectrometer was used for the proton identification, providing a $> 20 \text{ ns}$ separation from kaons and pions. The trigger efficiency of both spectrometer arms is at the 99.9% level and comes with a $\pm 0.1\%$ uncertainty. The correction for the proton absorption in the spectrometer comes with a small uncertainty of 0.18%. The tracks are traced, through the spectrometer optics, to the target to provide the particle momentum, scattering angle and target position information. Both spectrometers offer a better than 0.1% momentum resolution and an angular resolution of $\sim 1 \text{ mrad}$. The determination of the scattering angle comes with a 0.5 mrad uncertainty.

The coincidence time was determined as the difference in the time-of-flight between the two spectrometers, accounting for path-length variation corrections from the central trajectory and for the individual start-times. The experimental setup offered a better than 1 ns (FWHM) resolution in the coincidence timing spectrum that was measured within an 80 ns timing window. The random coincidences were subtracted using the side (accidental) bands of the coincidence time spectrum. The live-time correction, that accounts for the electronics and computer dead-time, came with an uncertainty that ranged between 0.3% and 0.6% for the different kinematical settings of the experiment.

The events of the exclusive reaction $ep \rightarrow ep\gamma$ were identified from the missing-mass reconstruction, through a selection cut around the photon peak in the missing-mass-squared spectrum. The contamination from the missing mass tail of the pion electroproduction events was studied in two methods. The contributions were evaluated by pion electroproduction simulation studies that employ the well known cross section of the reaction and offer an accurate description of the missing mass tail. In a second method, the pion contamination was determined through a phenomenological parametrization of the missing mass spectrum. The two methods exhibit a sub-percent agreement in the extracted cross section. A 1% was assigned as an uncertainty to this correction.

Elastic scattering measurements with a proton target were performed at different stages of the experiment, for calibration and normalization studies. A real-time normalization cross check during the measurement of the VCS cross section

was also performed from the simultaneous measurement of the $ep \rightarrow ep\pi^0$ reaction. In both the elastic and pion electro-production measurements, we found an excellent agreement to these well known cross sections, and confirmed that the spectrometer acceptance is accurately described by the simulation of the experiment.

The true momentum settings of the two spectrometers were determined based on a cross-calibration method that utilizes the pair of the azimuthal asymmetry measurements. Here, the momentum and position of the electron spectrometer remain the same between the two kinematical settings. The momentum setting for the proton spectrometer also remains constant, while the proton spectrometer is repositioned symmetrically with respect to the momentum transfer direction. Since the two kinematical settings involve identical momentum settings for each of the two spectrometers, the determination of their true momentum settings comes from a unique solution for both kinematics, that simultaneously calibrates the reconstructed VCS and pion electroproduction missing mass peaks to the true physical mass values for the photon and the pion, respectively. The correction between the set and the true values in the central momentum of the two spectrometers was smaller than 0.1%.

Cross sections and generalized polarizabilities. The measurement of the absolute VCS cross section, $\sigma \equiv d^5\sigma/dE'_e d\Omega'_e d\Omega_{cm}$, requires the determination of the five-fold solid angle, where dE'_e , $d\Omega'_e$ is the differential energy and solid angle of the scattered electron in the laboratory frame and $d\Omega_{cm}$ is the differential solid angle of the proton in the center-of-mass. The determination of the five-fold solid angle is calculated by using the Monte Carlo simulation program, SIMC. The simulation integrates a realistic description of the beam configuration, target geometry, spectrometer acceptances, resolution effects, energy losses and the radiative corrections as described in⁴⁵. The measured cross section is derived as:

$$\sigma = \frac{N}{\epsilon_{LT} \cdot \epsilon_{trk} \cdot \epsilon_{trig} \cdot \epsilon_p \cdot L \cdot \Delta\Omega^5} f_{rad} \frac{\sigma_P^{sim}}{\sigma_{avg}^{sim}} \quad (3)$$

The parameters of Eq. 3 are as follows: N is the number of the measured events within the acceptance cuts after the subtraction of the contributions arising from the target walls and from the accidental background, $\Delta\Omega^5$ is the five-fold coincidence solid angle that is determined from the simulation of the experiment, ϵ_{LT} denotes the efficiency that is associated with the computer and electronics live-time, ϵ_{trk} and ϵ_{trig} are the combined tracking and trigger efficiencies for the two spectrometers, respectively and ϵ_p is the efficiency that corrects for the proton absorption in the spectrometer. The f_{rad} denotes the radiative corrections and the luminosity $L = \frac{\rho * l * N_A}{A} * \frac{Q}{e}$, where ρ is the target density in g/cm^3 , l is the target length in cm , N_A is Avogadro's number, A is the mass number, Q is the accumulated charge of the measurement and e is the charge of electron. The σ_P^{sim} denotes the point cross section of the simulation at the central kinematics of each

bin, while the σ_{avg}^{sim} is the average cross section of the simulation within the analysis bin. The term $\sigma_P^{sim} / \sigma_{avg}^{sim}$ effectively provides the bin-centering correction for the extraction of the point cross section from the finite phase space of the analysis bin. A first-layer global phase-space cut in the data analysis selects the central 1/2 part of the coincidence acceptance, so that any potential influence from acceptance edge-effects is eliminated. The bin width size ($Q^2, W, \theta_{\gamma^* \gamma}, \phi_{\gamma^* \gamma}$) is varied in the analysis so as to validate the stability of the results as a function of the bin-size selection, and to confirm the good understanding of the coincidence phase space in the experiment simulation. The generated events in the simulation are weighted with a cross section using the DR model for VCS^{14–16}. In the DR formalism, the two scalar GPs come as an unconstrained part and can be adjusted as free parameters, while the proton electromagnetic form factors are introduced as an input. For the non-Born VCS part, a realistic initial parametrization is applied based on the current knowledge of the GPs. We extract the GPs from the measured cross sections through a fit that employs the dispersion relations model. The experimental cross sections are compared to the DR model predictions for all possible values for the two GPs, and the $\alpha_E(Q^2)$ and $\beta_M(Q^2)$ are fitted by a χ^2 minimization. Resolution and experimental parameters are studied by varying them in the analysis within their quoted precision, and their effect to the measured cross sections and to the extracted GPs is quantified as a systematic uncertainty. Other sources of systematic uncertainties involve the radiative corrections⁴⁵ that introduce a 1.5% uncertainty to the measured cross section, and the uncertainty in the determination of the coincidence solid angle that is better than 1.5%. The bin-centering correction was studied by varying the cross section model in the simulation and was found to be very small. We also adopt an additional 0.5% of uncertainty to account for any potentially undetermined small sources of systematic uncertainty in the measured cross section. The cross section results are reported in the Extended Data Tables 1, 2 and 3. The extracted generalized polarizabilities are reported in the Extended Data Table 4.

Induced polarization. We derive the transverse position space dependence of the induced polarization in the proton following³⁰:

$$\vec{P}_0(\vec{b}) = \hat{b} \int_0^\infty \frac{dQ}{(2\pi)} Q J_1(bQ) A(Q^2), \quad (4)$$

where \vec{b} is the transverse position, $b = |\vec{b}|$, $\hat{b} = \vec{b}/b$ and J_1 the 1st order Bessel function. The A is a function of the GPs:

$$A = -(2M) \sqrt{\tau} \sqrt{\frac{3}{2}} \sqrt{\frac{1+2\tau}{1+\tau}} \times \left\{ -P^{(L1,L1)0} + \frac{1}{2} P^{(M1,M1)0} - \sqrt{\frac{3}{2}} P^{(L1,L1)1} - \sqrt{\frac{3}{2}} (1+\tau) \left[P^{(M1,M1)1} + \sqrt{2} (2M\tau) P^{(L1,M2)1} \right] \right\}. \quad (5)$$

The GPs are expressed in the multipole notation $P^{(\rho' l', \rho l) S}$ ⁴⁶, where ρ (ρ') refers to the Coulomb/electric (L), or magnetic (M) nature of the initial (final) photon, l ($l' = 1$) is the angular momentum of the initial (final) photon, and S differentiates between the spin-flip ($S = 1$) and non spin-flip ($S = 0$) transition at the nucleon side. The $\tau \equiv Q^2/(4M^2)$, with M the nucleon mass. The two scalar GPs are defined as:

$$\alpha_E(Q^2) = -\frac{e^2}{4\pi} \sqrt{\frac{3}{2}} P^{(L1, L1)0}(Q^2) \quad (6)$$

$$\beta_M(Q^2) = -\frac{e^2}{4\pi} \sqrt{\frac{3}{8}} P^{(M1, M1)0}(Q^2) \quad (7)$$

with $e^2/4\pi = 1/\alpha_{QED} = 1/137$. In calculating the A function, the spin GPs are fixed by the dispersion relations^{15,16}. For the asymptotic part of $\alpha_E(Q^2)$ we use the parametrization that we derive from a fit to the world data:

$$\alpha_E(Q^2) = p0 * e^{-0.5 * (\frac{Q^2 - p1}{p2})^2} + \frac{1}{(p3 + Q^2/p4)^2} (fm^3) \quad (8)$$

with $p0 = (17.3 \pm 7.1)10^{-5}$, $p1 = 0.353 \pm 0.001$, $p2 = 0.040 \pm 0.021$, $p3 = 34.185 \pm 1.120$ and $p4 = 0.015 \pm 0$ (see "Experimental Fit" in Fig. 4(a)). For $\beta_M(Q^2)$ we find that the world data are described accurately by the DR model^{15,16} that adopts a single dipole behavior for the unconstrained part of the scalar GPs with a mass scale parameter of $\Lambda_\beta = 0.7 GeV$, and we adopt this parametrization.

Electric and magnetic polarizability radius. The electric polarizability radius $\langle r_{\alpha_E}^2 \rangle$ is extracted from Eq. 2. For $\alpha_E(0)$ we adopt the most recent measurement from³¹. In order to determine the slope of the electric GP at $Q^2 = 0$ we explore a variety of functional forms, namely combinations of polynomial, dipole, gaussian and exponential functions. We determine those functional forms that can provide a good fit to the data and a meaningful extraction of the slope in terms of its uncertainty. The fits are explored in two groups: one over the full Q^2 range, and a second one within a limited range at low- Q^2 that does not include the α_E anomaly, namely for $Q^2 = [0, 0.28] GeV^2$. For the experiments where the polarizabilities have been derived by both the Dispersion Relations and the Low Energy Expansion analysis, the variance of the two results is treated as a model uncertainty for each data point. The data points from^{8,9} have not been included in the fits, considering the disagreement with the new measurements reported in this work. The results of the individual fits are shown in the Extended Data Fig. 2. The final value for $\langle r_{\alpha_E}^2 \rangle$ is determined from the weighted average of the results of the individual fits. The uncertainty of $\langle r_{\alpha_E}^2 \rangle$ receives contributions from two terms. One term involves the uncertainty of the weighted average. The second term is quantified from the weighted variance of the individual fit results and effectively reflects the model dependence on the choice of the fitted parametrization. This is similar to what has been followed in the past for the extraction of the proton charge radius from fits that employ multiple functional forms

e.g.^{40,41}. The new result for $\langle r_{\alpha_E}^2 \rangle$ updates the earlier extractions^{2,33} of this quantity, as shown in Extended Data Fig. 3. In comparison to these results, the past derivations of $\langle r_{\alpha_E}^2 \rangle$ ^{2,33} have been performed considering a fit of a single function within a limited Q^2 -range, and an older measurement for $\alpha_E(0)$.

The mean square magnetic polarizability radius is derived from the magnetic polarizability measurements following

$$\langle r_{\beta_M}^2 \rangle = \frac{-6}{\beta_M(0)} \cdot \frac{d}{dQ^2} \beta_M(Q^2) \Big|_{Q^2=0}. \quad (9)$$

For $\beta_M(0)$ we adopt the most recent measurement from³¹. In order to determine the slope of the magnetic GP at $Q^2 = 0$ we explore a variety of functional forms, namely combinations of polynomial, dipole and exponential functions. We determine those functional forms that can provide a good fit to the data and a meaningful extraction of the slope in terms of its uncertainty e.g. for the dipole and (dipole-polynomial) functions we find that they give a good fit, but the radius is derived with a very large uncertainty and does not influence the final extraction of this quantity. The results of the individual fits are shown in the Extended Data Fig. 4. The exponential fit employs only 2 free parameters and offers an uncertainty ($\langle r_{\beta_M}^2 \rangle = 0.41 \pm 0.10 fm^2$) that is significantly smaller compared to the other functional forms that involve 3 or more free parameters. The fitted exponential curve also appears to be systematically different compared to the rest of the fitted functions, as can be seen in the Extended Data Fig. 4. In order not to bias the final extraction of the radius by the small uncertainty (or equivalently, the large weight factor) of this one fit, the fitted results are divided into two groups, one for only the exponential fit and a second group for the rest of the functions. For the second group, we derive the radius based on the weighted average and the weighted variance of the individual fits and we find $\langle r_{\beta_M}^2 \rangle = 0.85 \pm 0.25 fm^2$. We adopt the mean average of the two group values as the final result for the magnetic polarizability radius. For the uncertainty, we consider the spread of the two group values as a model uncertainty and we add it linearly to the statistical uncertainty. We find that $\langle r_{\beta_M}^2 \rangle = 0.63 \pm 0.31 fm^2$.

References

44. Mkrtchyan, H. *et al.* The lead-glass electromagnetic calorimeters for the magnetic spectrometers in Hall C at Jefferson Lab. *Nucl. Instrum. Meth.* **A719**, 85 (2013).
45. Vanderhaeghen, M., Friedrich, J.M., Lhuillier, D., Marchand, D., Van Hoorebeke, L., and Van de Wiele, J. QED radiative corrections to virtual Compton scattering. *Phys. Rev. C* **62**, 025501 (2000).
46. Guichon, P., Liu, G., and Thomas, A. W. Virtual Compton scattering and generalized polarizabilities of the proton. *Nucl. Phys.* **A591**, 606-638 (1995).

Data availability

The raw data from the experiment are archived in Jefferson Laboratory's mass storage silo and at Temple University, Department of Physics. The filtered data are archived at Temple University. The data are available from the authors upon request.

Code availability

The data analysis uses the standard C++ ROOT framework, which was developed at CERN and is freely available at <https://root.cern.ch>. The simulation of the experiment was generated with the Jefferson Lab simulation code SIMC. The DR fit was done using the dispersion relations code developed by B. Pasquini^{14–16}. The computer codes used for the data analysis are available upon request.

Acknowledgements

We would like to thank M. Vanderhaeghen as this work received great benefit from his input and suggestions. This work has been supported by the US Department of Energy Office of Science, office of Nuclear Physics under contract no. DE-SC0016577.

Author contributions

N.S. is the spokesperson of the experiment. He initiated, guided and supervised this effort. R.L worked on the data analysis and on the extraction of the cross sections and generalized polarizabilities. H.A. worked on the data analysis and on the extraction of the electric polarizability radius and of the induced polarization density. M.K.J. and M.P. co-supervised the data analysis efforts and worked on the simulation of the experiment. The entire VCS collaboration participated in the data collection and in the on-line analysis of the experiment.

Competing interests

The authors declare no competing interests.

Supplementary information

Supplementary information is available for this paper in the form of Extended Data figures and tables.

Correspondence and request of materials

Correspondence and request of materials should be addressed to N.S.

Extended Data




Figure 1. Cross section measurements of the VCS reaction for out-of-plane kinematics

a) Cross section measurements for out-of-plane kinematics at $Q^2 = 0.28 \text{ GeV}^2$. Results are shown for different bins in the total c.m. energy of the (γp) system, W . Top, middle and bottom panels correspond to $\phi_{\gamma^* \gamma} = 140^\circ$, $\phi_{\gamma^* \gamma} = 28^\circ$ and $\phi_{\gamma^* \gamma} = 38^\circ$, respectively. **b)** Measurements for out-of-plane kinematics at $Q^2 = 0.33 \text{ GeV}^2$. Top and bottom panels correspond to $\phi_{\gamma^* \gamma} = 150^\circ$ and $\phi_{\gamma^* \gamma} = 30^\circ$, respectively. **c)** Measurements for out-of-plane kinematics at $Q^2 = 0.40 \text{ GeV}^2$. Top, middle and bottom panels correspond to $\phi_{\gamma^* \gamma} = 152^\circ$, $\phi_{\gamma^* \gamma} = 35^\circ$ and $\phi_{\gamma^* \gamma} = 50^\circ$, respectively. The solid curve shows the Dispersion Relations (DR) fit for the two scalar generalized polarizabilities. The dashed curve shows the Bethe-Heitler plus Born-VCS (BH+Born) cross section. The error bars correspond to the total uncertainty, at the 1σ or 68% confidence level.




Figure 2. Electric polarizability radius fits

Top panel: The mean square electric polarizability radius fits using combinations of different functional forms (exp, gaus, pol, dipole correspond to exponential, gaussian, polynomial and dipole functions, respectively). The fits denoted with solid lines were performed over the full Q^2 -range of the world-data. The three functional forms denoted with dashed lines were performed in the low- Q^2 range, namely in $Q^2 = [0, 0.28] \text{ GeV}^2$. Bottom panel: The extracted mean square electric polarizability radius from the individual fits. The error bars correspond to the total uncertainty, at the 1σ or 68% confidence level. The blue band marks the final value for the extracted $\langle r_{\alpha_E}^2 \rangle = 1.28 \pm 0.24 \text{ fm}^2$.




Figure 3. Electric polarizability radius

The electric polarizability radius $r_{\alpha_E} \equiv \sqrt{\langle r_{\alpha_E}^2 \rangle}$ derived from this work is compared to the previous extractions of this quantity (red open symbols). The recent measurements of the proton charge radius r_E (blue symbols) are also shown. The error bars correspond to the total uncertainty, at the 1σ or 68% confidence level.




Figure 4. Magnetic polarizability radius fits

Top panel: The mean square magnetic polarizability radius fits using combinations of different functional forms (exp, pol, dipole correspond to exponential, polynomial and dipole functions, respectively). Bottom panel: The extracted mean square magnetic polarizability radius from the individual fits. The error bars correspond to the total uncertainty, at the 1σ or 68% confidence level. The blue band marks the final value for the extracted $\langle r_{\beta_M}^2 \rangle = 0.63 \pm 0.31 \text{ fm}^2$.