

Reference List

- Antonelli, A., Baldini, R., Biagini, M. E., & Calcaterra, S. (1992). Measurement of the $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ and $e^+e^- \rightarrow \omega\pi^+\pi^-$ reactions in the energy interval 1350–2400 MeV. *Zeitschrift Für Physik C ...*, 56(1), 15–19. <http://doi.org/10.1007/BF01589702>
- Antonelli, A., Baldini, R., Calcaterra, S., & Capon, G. (1988). Measurement of the reaction $e^+e^- \rightarrow \eta\pi^+\pi^-$ in the center of mass energy interval 1350–2400 MeV. *Physics Letters B*, B212(1), 133–138. [http://doi.org/10.1016/0370-2693\(88\)91250-6](http://doi.org/10.1016/0370-2693(88)91250-6)
- Aubert, B., Barate, R., Boutigny, D., & al, E. (2003). Study of the $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ process using initial state radiation with BABAR. *Physical Review D*.
- Aubert, B., Barate, R., Boutigny, D., & al, E. (2004). ... $\pi^+\pi^-\pi^+\pi^-$, $K^+K^-\pi^+\pi^-$, and $K^+K^-\pi^+\pi^-$ cross sections at center-of-mass energies 0.5–4.5 GeV measured with initial-state radiation. *Physical Review D*.
- Aubert, B., Barate, R., Boutigny, D., & al, E. (2005a). Study of $e^+e^- \rightarrow \rho^0\rho^0$ using initial state radiation with BABAR. *Physical Review D*.
- Aubert, B., Barate, R., Boutigny, D., Couderc, F., Karyotakis, Y., Lees, J. P., et al. (2005b). $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$, $K^+K^-\pi^+\pi^-$, and $K^+K^-\pi^+\pi^-$ cross sections at center-of-mass energies 0.5–4.5 GeV measured with initial-state radiation. *Physical Review D*, 71(5), 052001. <http://doi.org/10.1103/PhysRevD.71.052001>
- Aubert, B., Bona, M., Boutigny, D., & al, E. (2007). Measurements of $e^+e^- \rightarrow K^+K^-\eta$, $K^+K^-\pi^0$, and $K^+K^-\pi^0$ cross sections using initial state radiation events. *Physical Review D*.
- Bisello, D., Bizot, J. C., Buon, J., Cordier, A., Delcourt, B., others. (1981). Study of the Reaction $e^+e^- \rightarrow 3\pi^+3\pi^-$ in the Total Energy Range 1400-MeV to 2180-MeV. *Phys.Lett.*, B107(1-2), 145–147. [http://doi.org/10.1016/0370-2693\(81\)91169-2](http://doi.org/10.1016/0370-2693(81)91169-2)
- Bisello, D., Busetto, G., Castro, A., Nigro, M., Penzo, M., Pescara, L., et al. (1988). Study of the reaction $e^+e^- \rightarrow K^+K^-$ in the energy range $1350 < \sqrt{s} < 2400$ MeV. *Zeitschrift Für Physik C Particles and Fields*, 39(1), 13–19. <http://doi.org/10.1007/BF01560386>
- Bisello, D., Busetto, G., Castro, A., Nigro, M., Pescara, L., Sartori, P., et al. (1991). Observation of an isoscalar vector meson at ≈ 1650 MeV/c² in the $e^+e^- \rightarrow K^+\bar{K}^0\pi^0$ reaction. *Zeitschrift Für Physik C Particles and Fields*, 52(2), 227–230. <http://doi.org/10.1007/BF01560440>
- Buon, J., Bisello, D., Bizot, J. C., Cordier, A., Delcourt, B., others. (1982). Interpretation of Dm1 Results on e^+e^- Annihilation Into Exclusive Channels Between 1.4-GeV and 1.9-GeV With a $\rho^0\omega\phi$ Model. *Phys.Lett.*, B118(1-3), 221–224. [http://doi.org/10.1016/0370-2693\(82\)90632-3](http://doi.org/10.1016/0370-2693(82)90632-3)
- Clegg, A. B., & Donnachie, A. (1994). Higher vector meson states produced in electron - positron annihilation. *Z.Phys.*, C62(3), 455–470. <http://doi.org/10.1007/bf01555905>
- Collaboration, T. B., & Aubert, B. (2009, August 25). Precise measurement of the $e^+e^- \rightarrow \pi^+\pi^- (\gamma)$ cross section with the Initial State Radiation method at BABAR. *arXiv.org*. <http://doi.org/10.1103/PhysRevLett.103.231801>

Cordier, A., Bisello, D., Bizot, J. C., Buon, J., Delcourt, B., & Mané, F. (1981). Observation of a new isoscalar vector meson in $e^+e^- \rightarrow \omega\pi^+\pi^-$ annihilation at 1.65 GeV. *Physics Letters B*, 106(1-2), 155–158. [http://doi.org/10.1016/0370-2693\(81\)91101-1](http://doi.org/10.1016/0370-2693(81)91101-1)

Cordier, A., Bisello, D., Bizot, J. C., Buon, J., Delcourt, B., others. (1982). Study of the $e^+e^- \pi^+\pi^- \pi^+\pi^-$ Reaction in the 1.4-GeV to 2.18-GeV Energy Range. *Phys.Lett.*, B109(1-2), 129–132. [http://doi.org/10.1016/0370-2693\(82\)90478-6](http://doi.org/10.1016/0370-2693(82)90478-6)

Delcourt, B., Bisello, D., Bizot, J. C., Buon, J., Cordier, A., others. (1981). Study of the Reaction $e^+e^- K^+K^-$ in the Total Energy Range 1400-MeV to 2060-MeV. *Phys.Lett.*, B99(3), 257–260. [http://doi.org/10.1016/0370-2693\(81\)91121-7](http://doi.org/10.1016/0370-2693(81)91121-7)

Lees, J. P. (2011). Cross Sections for the Reactions $e^+e^- \rightarrow K^+K^-\pi^+\pi^-$, $K^+K^-\pi^0\pi^0$, and $K^+K^-\pi^+\pi^-$ Measured Using Initial-State Radiation, 1–35.

Lees, J. P., Poireau, V., Tisserand, V., Tico, J. G., Grauges, E., Martinelli, M., et al. (2012). Initial-State Radiation Measurement of the $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$ Cross Section. *Nuclear Instruments and Methods, hep-ex*.

Mané, F., Bisello, D., Bizot, J. C., Buon, J., Cordier, A., others. (1981). Study of the Reaction $e^+e^- K_S^0 K_L^0$ in the Total Energy Range 1.4-GeV to 2.18-GeV and Interpretation of the K^+ and K^0 Form-factors. *Phys.Lett.*, B99(3), 261–264. [http://doi.org/10.1016/0370-2693\(81\)91122-9](http://doi.org/10.1016/0370-2693(81)91122-9)