

# Search for a Dark Photon in Electro-Produced $e^+e^-$ Pairs with the Heavy Photon Search Experiment at JLab

---

**Omar Moreno<sup>a\*</sup>, Matthew Solt<sup>b\*</sup>**

<sup>a</sup> SLAC National Accelerator Laboratory

<sup>b</sup> Stanford University

E-mail: [omoreno@slac.stanford.edu](mailto:omoreno@slac.stanford.edu), [mrsolt@slac.stanford.edu](mailto:mrsolt@slac.stanford.edu)

The Heavy Photon Search experiment took its first data in a 2015 engineering run at the Thomas Jefferson National Accelerator Facility, searching for an electro-produced dark photon. Using 1.7 days ( $1170 \text{ nb}^{-1}$ ) of data, a search for a resonance in the  $e^+e^-$  invariant mass distribution between 19 and 81 MeV/c<sup>2</sup> showed no evidence of dark photon decays above the large QED background, confirming earlier searches and demonstrating the full functionality of the experiment. Upper limits on the square of the coupling of the dark photon to the Standard Model photon are set at the level of  $6 \times 10^{-6}$ . In addition, a search for displaced dark photon decays did not rule out any territory but resulted in a reliable analysis procedure that will probe hitherto unexplored parameter with future, higher luminosity runs.

*XXXIX International Conference on High Energy Physics  
4-11 July 2018  
Seoul, South Korea*

---

\*Speaker

## 1. Introduction

There is compelling cosmological and astrophysical evidence from galactic rotation curves, gravitational lensing, and the Cosmic Microwave Background that  $\sim 85\%$  of the total matter in the universe is made of invisible matter. The particles in the Standard Model cannot account for this invisible matter, or dark matter. Currently, Weakly Interacting Massive Particles (WIMPs) are the most popular model of dark matter due to its motivation of the so-called WIMP Miracle where the thermal production of dark matter in the early universe can successfully achieve the observed relic abundance of dark matter with a new particle with a mass and coupling close to the weak-sector scale. In addition, Supersymmetry (SUSY) described the same type of object. However, after decades of searching for WIMPs, many dark matter detection experiments have failed to detect a WIMP and experiments at the Large Hadron Collider have failed to detect SUSY particles, and parameter space is quickly running out of room for the simplest versions of these models.

This leaves room for other more novel dark matter scenarios. One such model called hidden sectors proposes a whole zoo of new particles that do not interact directly with Standard Model particles. In order for these hidden sector particles to be a candidate for thermal dark matter; however, there must be an indirect interaction between a hidden sector particle and Standard Model particles.

A heavy photon (or dark photon or  $A'$ ) is a natural candidate for such an interaction. A heavy photon would be a mediator of a hidden  $U'(1)$  gauge symmetry that interacts with SM particles through kinetic mixing of the  $U(1)_Y$  (hypercharge) gauge boson. This kinetic mixing generates an interaction between the SM photon and the heavy photon which allows an indirect coupling of the heavy photon to electric charge.

Because of this effective coupling of the heavy photon to electric charge, heavy photons can be produced in a process analogous to bremsstrahlung radiation and then decay to charged lepton pairs. The Heavy Photon Search Experiment is a fixed target experiment that utilizes this mechanism by producing heavy photons from a high energy electron beam incident on a thin tungsten target and then detecting the  $e^+e^-$  decay products. HPS can explore the  $A'$  mass range of  $18 \text{ MeV}/c^2$  to  $500 \text{ MeV}/c^2$ . HPS distinguishes these decay products from radiative and Bethe-Heitler trident backgrounds that also have  $e^+e^-$  final states through two methods. The first is a bump hunt for an  $A'$  with large coupling in which a resonance peak at the  $A'$  mass over a large QED background. For a sufficiently small  $A'$  coupling, a second analysis can be performed by searching for a displaced vertex signature (in the range 1-10 cm) past a background of prompt QED tridents.

The HPS detector was installed and commissioned in the spring of 2015 in the Hall B alcove at Jefferson Laboratory followed by a short engineering run. These proceedings report both the resonance and displaced vertex searches for the 2015 engineering run in which a total of  $1165 \text{ nb}^{-1}$  (7.25 mC of charge) were collected at a beam energy of 1.056 GeV and beam current of 50 nA.


## 2. Detector Overview

The HPS detector is a compact, large forward spectrometer with high rate front end electronics located in the Hall B alcove at Jefferson Laboratory directly downstream of the CLAS-12 detec-

tor. The high luminosity electron beam is produced at the Continuous Electron Beam Accelerator Facility (CEBAF) at Jefferson Laboratory and is incident on a thin tungsten target for  $A'$  production.

Since an  $A'$  takes most of the incident beam energy, the  $e^+e^-$  decay products are highly boosted and the design of the HPS detector must have excellent forward acceptance. However, the forward acceptance is limited by occupancies due to a high rate of beam electrons. To account for this, the detector is split into two halves (top and bottom) to allow the majority of beam electrons to avoid interaction with the detector. In addition, the HPS detector must also have excellent momentum, vertex, and mass resolutions for both a resonance search and a displaced vertex search.

The HPS detector consists of three major components shown in Figure 1 - a large dipole magnet, a silicon vertex tracker (SVT), and a highly segmented lead tungstate electromagnetic calorimeter (Ecal). Together, these subsystems provide complete kinematic information. The dipole magnet provides curvature for momentum measurement and particle identification. For 2015 running, the magnetic field strength was 0.25 T. The Ecal is positioned at the back of the detector and is used as the primary trigger and for particle identification.



**Figure 1:** Schematic of the Heavy Photon Search Detector used during the 2015 engineering run.

The SVT is positioned at the heart of the detector and is used to track particle trajectories for vertex reconstruction. The SVT consists of six measurement layers of silicon microstrips on both top and bottom, and each layer consists of two silicon sensors where one is oriented at a small stereo angle (50 or 100 mrad) in order to enable a full 3D hit reconstruction. The entirety of the SVT lies in a vacuum so that beam gas interactions are sufficiently minimized. The first silicon layer is located 0.5 mm from the beam plane for the best possible forward acceptance at 15 mrad, and is positioned about 10 cm downstream of a thin (0.125%  $X_0$ ) tungsten foil target for the best possible vertex resolution. The silicon sensors are thin (250 microns) in order to reduce the material budget in the SVT, and hence reduce multiple scattering.

During the 2015 engineering run, the HPS detector performed as expected with momentum resolution, mass resolution, and vertex resolutions all in agreement with predicted values.

### 3. Resonance Search

Searching for a heavy photon resonance requires accurate reconstruction of the  $e^+e^-$  invariant mass spectrum; rejection of background events due to converted WAB events, non-radiative tridents

from the Bethe-Heitler process, and occasional accidental  $e^+e^-$  pairs; and efficient selection of  $A'$  candidates. Selecting  $A'$  candidates is equivalent to selecting radiative tridents since they have identical kinematics for a given mass. In order to perform a blind search, the event selection was optimized using  $\sim 10\%$  of the 2015 engineering run dataset.

Heavy photon candidates are created from pairs of electron and positron tracks, one in each half of the SVT, each of which point to an energy cluster in the ECal. Each track must pass loose quality requirements and have a reconstructed momentum less than 75% of the beam energy ( $0.788 \text{ GeV}/c^2$ ) to reject scattered beam electrons. The background from accidental pairs was reduced to less than 1% by requiring the time between the ECal clusters be less than 2 ns and the time between a track and the corresponding cluster be less than 5.8 ns.

Trident production, comprised of radiative and Bethe-Heitler processes, is the main physics background to HPS. Although the Bethe-Heitler diagram dominates over the radiative process at all pair momenta, it is kinematically distinct. As a result, the contribution of the Bethe-Heitler process to the final event sample can be minimized by requiring the momentum sum of the  $e^+e^-$  pair to be greater than 80% of the beam energy ( $0.84 \text{ GeV}/c^2$ ), where the radiative tridents are peaked.

The other significant source of background arises from converted WAB events in which the bremsstrahlung photon is emitted at a large angle ( $> 15 \text{ mrad}$ ), converts in the target, first or second layer of the SVT, and gives rise to a detected positron in the opposite half of the detector from the recoiling incoming electron. The converted WAB background was substantially reduced by requiring that both tracks have hits in the first two layers of the SVT. Furthermore, applying additional requirements on the transverse momentum asymmetry between the  $e^+e^-$  ( $< .47$ ) and the transverse distance of closest approach to the beam spot of the positron track ( $< 1.1 \text{ mm}$ ) reduces the contamination from converted WABs down to 12%.

A prompt heavy photon is expected to appear as a Gaussian-shaped resonance above the  $e^+e^-$  invariant mass spectrum, centered on the  $A'$  mass and with a width,  $\sigma_{m_{A'}}$ , which characterizes the experimental mass resolution. Calibration of the  $A'$  mass scale and resolution is done by using Møller scattering ( $e^-e^- \rightarrow e^-e^-$ ) events. The resulting parametrization is shown in Figure ?? and is used as an input to the resonance search.


Since the mass of the putative  $A'$  is unknown a priori, the entire  $e^+e^-$  invariant mass spectrum is scanned for any significant peaks. This search is performed in a broad mass window around each candidate mass, repeated in 0.5 MeV steps between 19 and 81 MeV. Within the window, the distribution of  $A'$  signal events is modeled using the probability distribution function

$$P(m_{e^+e^-}) = \mu \cdot \phi(m_{e^+e^-} | m_{A'}, \sigma_{m_{A'}}) + B \cdot \exp(p(m_{e^+e^-} | \mathbf{t})) \quad (3.1)$$

where  $m_{e^+e^-}$  is the  $e^+e^-$  invariant mass,  $\mu$  is the signal yield,  $B$  is the number of background events within the window,  $\phi(m_{e^+e^-} | m_{A'}, \sigma_{m_{A'}})$  is a Gaussian probability distribution describing the signal and  $p(m_{e^+e^-} | \mathbf{t})$  is a Chebyshev polynomial of the first kind with coefficients  $\mathbf{t} = (t_1, \dots, t_j)$  that is used to describe the background shape. From optimization studies, a 5th (3rd) order Chebyshev polynomial was found to best describe the background below (above) 39 MeV. Note that  $m_{A'}$  and  $\sigma_{m_{A'}}$  are set to the  $A'$  mass hypothesis and experimental mass resolution, respectively. Estimating the signal yield, the background normalization, and the background shape parameters within a window is done with a binned maximum likelihood fit using a bin width of 0.05 MeV. A detailed

discussion of the procedures followed can be found in [?]. Briefly, the log of the ratio of likelihoods for the background-only fit to that of the best signal-plus-background fit provides a test statistic from which the  $p$ -value can be calculated, giving the probability that the observed signal is a statistical fluctuation. The  $p$ -value is corrected for the “Look Elsewhere Effect” (LEE) by performing simulated resonance searches on 4,000 pseudo data sets [?].

A search for a resonance in the  $e^+e^-$  invariant mass spectrum found no evidence of an  $A'$  signal. The most significant signal was observed at 37.7 MeV and has a local  $p$ -value of 0.17%. After accounting for the LEE correction, the most significant  $p$ -value is found to have a global  $p$ -value of 17% corresponding to less than  $2\sigma$  in significance. Since no significant signals were found, a 95% C.L. upper limit is set, power-constrained [?] to the expected limit.



**Figure 2:** The 95% C.L. power-constrained [?] upper limits on  $\epsilon^2$  versus  $A'$  mass obtained in this analysis. A limit at the level of  $6 \times 10^{-6}$  is set. Existing limits from beam dump [?, ?, ?, ?, ?, ?, ?, ?], collider [?, ?, ?, ?, ?] and fixed target experiments [?, ?, ?, ?] are also shown. The region labeled “ $a_e$ ” is an exclusion based on the electron  $g - 2$  [?, ?, ?, ?]. The green band labeled “ $a_\mu \pm 2\sigma$ ” represents the region that an  $A'$  can be used to explain the discrepancy between the measured and calculated muon anomalous magnetic moment [?, ?].

The proportionality between  $A'$  and radiative trident production allows the normalization of the  $A'$  rate to the measured rate of trident production [?]. This leads to a relation that allows the signal upper limit,  $S_{up}$ , to be related to the  $A'$  coupling strength as

$$\epsilon^2 = \left( \frac{S_{up}/m_{A'}}{f\Delta B/\Delta m} \right) \left( \frac{2N_{eff}\alpha}{3\pi} \right) \quad (3.2)$$

where  $N_{eff}$  is the number of decay channels kinematically accessible (=1 for HPS searches below the dimuon threshold),  $\Delta B/\Delta m$  is the number of background events per MeV,  $\alpha$  is the fine structure constant and  $f = 8.5\%$  is the fraction of radiative trident events comprising the background. Using equation 3.2, the limits on  $\epsilon$  set by HPS are shown on Figure 2.

The reach shown in Figure 2 includes all statistical and systematic uncertainties. The main systematic uncertainties on the signal yields arise from the uncertainty in the mass resolution (3%)

and biases observed in the fit due to the background and signal parameterization (1.3-1.5%, depending on mass). When scaling the extracted signal yield upper limits to a limit on  $\epsilon$ , the primary systematic uncertainty in the radiative fraction is due to the unknown composition of the final  $e^+e^-$  sample (7%). Many other possible sources of systematic uncertainty were investigated and accounted for but contribute negligibly to the result.

#### 4. Displaced Vertex Search


Searching for a long-lived heavy photon requires precise reconstruction of the vertex position of the  $e^+e^-$  pair, rejection of background events that reconstruct significantly downstream of the target, and sufficient signal efficiency for a displaced  $A'$ . The precision of the vertex position is limited by multiple scattering in the SVT, particularly the first layer. The event selection is optimized for choosing well-reconstructed events (good tracks and vertex reconstruction) and tries to eliminate events which arise from scatters that could produce downstream reconstructed vertices. From  $A'$  simulation, cuts that improve our selection of events that have favorable kinematics were implemented for selecting heavy photons.

Much of the initial event selection is similar to the resonance search in Section 3 such as timing cuts on tracks and Ecal clusters, track quality requirements, requiring an  $e^+e^-$  pair to be in opposite halves of the SVT, requiring tracks to be matched to ECal clusters, rejection of scattered beam electrons and WABs, and the radiative cut to select  $e^+e^-$  pairs with energy sum near the beam energy. Specifically for the vertexing analysis, excellent vertex quality (based on the DOCA of the  $e^+$  and  $e^-$  tracks) with total momentum that points back to the beam spot is required. In order to reduce the effects of layer 1 mis-hits which could reconstruct a false vertex downstream of the target, an isolation cut that compares the distance between first layer hits and the impact parameter at the target was implemented and tracks that share 5 hits with another track were removed.

In addition, both tracks to having layer 1 and layer 2 SVT hits were required. The layer 1 requirement removes tracks that have degraded vertex resolution and particles that may have scattered in the layer 1 dead silicon (which naturally reconstruct a false vertex downstream of the target). The layer 1 requirement also has a significant effect on the reconstruction efficiency for long-lived, low-mass heavy photons as shown in the  $A'$  distribution in Figure 3.

The heavy photon production rate (cross section), for both prompt and displaced vertices, is proportional to the radiative trident cross section. In looking for heavy photons with displaced vertices, the vertex distribution is examined as a function of the reconstructed invariant mass of the  $e^+e^-$  pair as shown in Figure 3, and events of interest are identified as originating far beyond the tails of the prompt trident backgrounds.

When searching for a displaced heavy photon, a downstream region having little background must be selected. Therefore, a downstream  $z$  vertex position beyond which there should be fewer than 0.5 background events per mass bin, called a  $z_{cut}$ , was chosen. The  $z_{cut}$  varies as a function of mass as shown in red in Figure 3 and, ideally, can be selected to minimize backgrounds whilst maximizing  $A'$  reach efficiency. Based on Poisson statistics alone, the 90% confidence limit for zero background requires us to have an expected number of  $A'$  events greater than 2.3.



**Figure 3:** Left: The vertex distribution for a mass slice from data (blue) with all cuts applied and with Gaussian core fit (magenta) and downstream exponential tail fit (red) overlaid with an simulated  $A'$  Monte Carlo at the same mass (black). The tail distribution of the  $A'$  falls off more slowly compared to the distribution of events and scattering tails from the target but falls off rapidly beyond 40 mm downstream due to acceptance effects of requiring hits in Layer 1. For this particular mass slice, the  $z_{cut}$  was found to be at 37.6 mm. Right: The reconstructed  $z$  vertex is shown versus the reconstructed mass of the  $e^+e^-$  pair for all events in the 2015 L1L1 data set. The  $z_{cut}$  is obtained by slicing this distribution by mass hypothesis (window size of  $\pm 1.9\sigma_m$ ) and fitting each  $z$  vertex distribution with Eq. eq:vtxFit. The  $z_{cut}$  shown in red is the point beyond which 0.5 background events is expected based on the fit to the tail.

An upper limit on the heavy photon production at a given  $m_{A'}$  and  $\epsilon^2$  is the maximum rate at which heavy photons could be produced, and still be consistent with the data. The confidence level used for this analysis is 90%: in other words, if a heavy photon signal does exist at a given rate, the limit set by this analysis will (incorrectly) exclude that signal rate only 10% of the time.


The method chosen for setting limits is the “optimum interval” method by Yellin [S. Yellin, 2002]. This method was developed for dark matter direct detection experiments, and is intended for experiments where the signal shape is known, but the backgrounds are not fully understood and there is the possibility of an unexpected background. A particular strength of the method is that it minimizes the influence of a background that is concentrated in one part of the data distribution.

The optimum interval method sets a one-sided upper limit (with confidence level  $C$ ) on the number of signal events  $\mu$  in a one-dimensional data set, where the shape of the signal distribution is known. For HPS, the data set is the distribution of vertex  $z$  locations, after applying the mass (in mass bins proportional to the mass resolution) and  $z_{cut}$  cuts; the signal shape is the distribution reported in Figure 3 for the  $m_{A'}$  and  $\epsilon^2$  being tested. The maximum number of detectable  $A'$ s and the corresponding limit as determined by the optimum interval method is shown in Fig. 4.

By definition of  $z_{cut}$ , about 0.5 background events past  $z_{cut}$  per mass slice (about 4 total) are expected; however, Figure 3 show many more such events past  $z_{cut}$ . This indicates that our background model is not complete in describing the background events. Detailed Monte Carlo studies reveal a similar deviation from the background model with some events due to mis-hits



in layer 1 (which the isolation cut fails to eliminate) and large scatters due to a single Coulomb interaction (as opposed to multiple scattering) in the layer 1 silicon. Further work will need to be done to mitigate these backgrounds and better model them, which will be incorporated in future analysis.



**Figure 4:** Left: The number of detectable  $A'$  events after applying all the analysis cuts and integrating the signal shape in Figure 3 over  $z > z_{cut}$ . The number of detectable  $A'$ s is found to be a maximum of 0.097 events where  $A'$  production is maximal at a mass of 43.6 MeV and  $\epsilon^2$  coupling of  $2.4E - 9$ . Right: 90% CL upper limit on  $\mu/\mu_{exp}$ , the ratio of the true production rate to the expected production rate for a heavy photon. A value of 1 would mean exclusion; the lowest contour on this plot is 35.7 (which can be interpreted as just beginning to exclude an  $A'$ -like model with 35.7 times the cross section) at a mass of 51.4 MeV and coupling of  $1.7E - 9$ . The vertical ridges in this plot correspond to the locations of events in mass space.

## 5. Conclusion

Both a resonance search and a displaced vertex search for a heavy photon with a mass between 19 and 81 MeV which decays to an  $e^+e^-$  pair was performed. A search for a resonance in the  $e^+e^-$  invariant mass spectrum yielded no significant excess and established upper limits on the square of the coupling at the level of  $6 \times 10^{-6}$ , confirming results of earlier searches. A search for displaced vertices has been established, although no heavy photon parameter space could be excluded. While not covering new territory in both searches in this short engineering run, these searches did establish that HPS operates as designed and will, with future running, extend coverage for  $\epsilon^2$  below the level of  $10^{-6}$  in the resonance search and provide coverage of unexplored parameter space at smaller values of the coupling from a search for displaced vertices.

## 6. Acknowledgments

The authors are grateful for the outstanding efforts of the Jefferson Laboratory Accelerator Division and the Hall B engineering group in support of HPS. The research reported here is supported by the U.S. Department of Energy Office of Science, Office of Nuclear Physics, Office of High Energy Physics, the French Centre National de la Recherche Scientifique, United Kingdom's Science and Technology Facilities Council (STFC), the Sesame project HPS@JLab funded by the French region Ile-de-France and the Italian Istituto Nazionale di Fisica Nucleare. Jefferson Science Associates, LLC, operates the Thomas Jefferson National Accelerator Facility for the United States Department of Energy under Contract No. DE-AC05-06OR23177.



## References

- [P. Billoir, 1992] P. Billoir and S. Qian (1992). Fast Vertex Fitting with a Local Parametrization of Tracks  
[NIM Volume 311; Issues 1-2](#)
- [S. Yellin, 2002] S. Yellin (2002). Finding an Upper Limit in the Presence of Unknown Background.  
[arXiv:physics/0203002](#)
- [S. Yellin, 2011] S. Yellin (2011). Some Ways of Combining Optimum Interval Upper Limits.  
[arXiv:1105.2928](#)
- [O. Moreno, 2017] O. Moreno, N. Baltzell, M. Graham, and J. Jaros (2017). Search for a Heavy Photon in Electro-Produced  $e^+e^-$  Pairs with the HPS Experiment at JLab. [Analysis Note](#)
- [S. Uemera, 2016] S. Uemera (2016). Searching for Heavy Photons in the HPS Experiment. [PhD Thesis](#)
- [H. Szumila-Vance, 2017] H. Szumila-Vance (2017). Search for Heavy Photons with Detached Vertices in the Heavy Photon Search Experiment. [PhD Thesis](#)
- [H. Szumila-Vance, 2015] H. Szumila-Vance (2015). HPS Ecal Timing Calibration for the Spring 2015 Engineering Run. [HPS Note](#)
- [Heavy Photon Search Collaboration] HPS Collab. HPS Collaboration Confluence Site. [HPS Site](#)
- [Jefferson Lab] Jefferson Lab. Jefferson Lab Site. [Jefferson Lab Site](#)