

insert space [1]

hyphens to be consistent with other parts of the document [2]

0.1 Update on JEF experiment

The $\eta^{(\prime)}$ meson, with the quantum numbers of the vacuum, provides a unique, flavor-conserving laboratory to probe the isospin-violating sector of low-energy QCD and to search for new physics Beyond the Standard Model (BSM). The JEF experiment [1, 2] was approved by JLab Program Advisory Committee (PAC) in 2017 to perform precision measurements of various $\eta^{(\prime)}$ decays with emphasis on rare neutral modes, running in parallel with the GlueX experiment using an upgraded Forward Calorimeter (FCAL-II). Significantly boosted $\eta^{(\prime)}$ will be produced through $\gamma + p \rightarrow \eta^{(\prime)} + p$ with an 8-12 GeV tagged photon beam. The $\eta^{(\prime)}$ decay photons (in leptons) will be measured in an upgraded forward calorimeter with high-granularity, high-resolution PbWO₄ core in the central region that minimizes shower overlaps and optimizes invariant mass reconstruction. Non-coplanar backgrounds will be suppressed by tagging $\eta^{(\prime)}$ with recoil proton detection. Compared to the previous or planned η/η' experiments, such as A2-MAMI [3, 4], WASA-at-COSY [5], KLOE-II [7], BESIII [6] and the proposed future REDTOP [8], JEF is the only one producing highly boosted $\eta^{(\prime)}$ so that its detection efficiency is insensitive to the detector thresholds, thus less prone to the experimental systematics. The capability of tagging every $\eta^{(\prime)}$ in combination with the upgraded calorimeter offers two orders of magnitude improvement in background suppression. JEF will simultaneously produce η and η' at the similar rate, $\sim 6 \times 10^7$ tagged η and $\sim 5 \times 10^7$ tagged η' (with detected recoil protons) per 100 days of beam time. These make JEF a unique η and η' factory in the world with no competition in rare neutral decay modes.

Though the JEF experiment will offer sensitive probes for a broad range of physics topics as described in [1, 2], its primary objectives will focus on the following:

- Search for new sub-GeV gauge bosons.

Vector:

- A leptophobic vector boson (B') coupled to baryon number.

$$\begin{aligned} \eta, \eta' &\rightarrow B' \gamma \rightarrow \pi^0 \gamma \gamma, \quad (0.14 < m_{B'} < 0.62 \text{ GeV}) \\ \eta' &\rightarrow B' \gamma \rightarrow \pi^+ \pi^- \pi^0 \gamma, \quad (0.62 < m_{B'} < 1 \text{ GeV}) \end{aligned}$$

- A dark photon coupled to the Standard model photon via kinetic mixing [10, 11, 12, 13].

$$\eta, \eta' \rightarrow A' \gamma \rightarrow e^+ e^- \gamma$$

Scalar: Search for hadrophilic scalar.

$$\begin{aligned} \eta &\rightarrow \pi^0 S \rightarrow \pi^0 \gamma \gamma, \pi^0 e^+ e^-, \quad (10 \text{ MeV} < m_S < 2m_\pi) \\ \eta, \eta' &\rightarrow \pi^0 S \rightarrow 3\pi, \eta' \rightarrow \eta S \rightarrow \eta \pi \pi, \quad (m_S > 2m_\pi) \end{aligned}$$

Axion-Like Particle (ALP): Light pseudoscalars will be searched via

$$\eta, \eta' \rightarrow \pi \pi a \rightarrow \pi \pi \gamma \gamma, \pi \pi e^+ e^-$$

- A search for C-violating $\eta^{(\prime)}$ decays (such as $\eta^{(\prime)} \rightarrow 3\gamma$ and $\eta^{(\prime)} \rightarrow 2\pi^0\gamma$) and a mirror asymmetry in the Dalitz distribution of $\eta^{(\prime)} \rightarrow \pi^+\pi^-\pi^0$ will offer the best direct constraints on new C-violating, P-conserving reactions (CVPC). As pointed out by M. Ramsey-Musolf, the Electric Dipole Moment (EDM) searches may place no constraint on CVPC in the presence of a conspiracy or new symmetry; only the direct searches of CVPC are unambiguous [18].

author in ref [18] so no need to mention author by name [17]

Summary of Comments on index-20.dvi

Page: 1

	Number: 1	Author: zisis	Subject: Callout	Date: 2020-06-01, 8:10:58 AM
	insert space			
	Number: 2	Author: zisis	Subject: Callout	Date: 2020-06-01, 8:17:43 AM
	hyphens to be consistent with other parts of the document			
	Number: 3	Author: zisis	Subject: Line	Date: 2020-06-01, 8:18:00 AM
	Number: 4	Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:12:43 AM
	-			
	Number: 5	Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:12:47 AM
	-			
	Number: 6	Author: zisis	Subject: Inserted Text	Date: 2020-06-01, 8:11:28 AM
	mesons			
	Number: 7	Author: zisis	Subject: Comment on Text	Date: 2020-06-01, 8:12:03 AM
	and and/or?			
	Number: 8	Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:12:18 AM
	FCAL-II			
	Number: 9	Author: zisis	Subject: Cross-Out	Date: 2020-06-01, 8:13:35 AM
	Number: 10	Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:13:45 AM
	-			
	Number: 11	Author: zisis	Subject: Comment on Text	Date: 2020-06-01, 8:14:32 AM
	redundant; tagged eta was defined above			
	Number: 12	Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:19:04 AM
	are			
	Number: 13	Author: zisis	Subject: Inserted Text	Date: 2020-06-01, 8:19:07 AM
	A			
	Number: 14	Author: zisis	Subject: Inserted Text	Date: 2020-06-01, 8:20:09 AM
	(vector, scalar and axion-like particle)			
	Number: 15	Author: zisis	Subject: Comment on Text	Date: 2020-06-01, 8:15:30 AM
	move references after scalar			
	Number: 16	Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:16:16 AM
	The			
	Number: 17	Author: zisis	Subject: Callout	Date: 2020-06-01, 8:16:50 AM
	author in ref [18] so no need to mention author by name			

3. Probing the low-energy QCD via precision measurements. A low-background measurement of the rare decay $\eta \rightarrow \pi^0 2\gamma$ provides a clean, rare window into $\mathcal{O}(p^6)$ in chiral perturbation theory [19]. This is the only known meson decay that proceeds via a polarizability-type mechanism. The precision in the Dalitz distribution will be sufficient for the first time to explore the role of scalar meson dynamics and its interplay with the vector meson dominance. The measurements of the transition form factor of η and η' via the $\eta^{(\prime)} \rightarrow e^+e^-\gamma$ decays offer important input to calculate hadronic light-by-light corrections to the muon $g - 2$ [20].
4. $\eta \rightarrow 3\pi$ promises an accurate determination of the quark mass ratio, $\mathcal{Q} = (m_s^2 - \hat{m}^2)/(m_d^2 - m_u^2)$ with $\hat{m} = (m_u + m_d)/2$. A recent dispersive analysis result yields $\mathcal{Q} = 22.1 \pm 0.7$ [21]. The statistical uncertainty due to fitting the Dalitz distribution of $\eta \rightarrow \pi^+\pi^-\pi^0$ (4.7M events) from KLOE-II [22] contributes ± 0.44 to the total error budget for the extracted \mathcal{Q} [21], accounting for one of the biggest uncertainties. The A2-MAMI result for the neutral decay $\eta \rightarrow 3\pi^0$ with 7M events [23] was also considered in [21], however, it imposes less constraint on \mathcal{Q} because of identical final state pions. JEF will be able to reduce this uncertainty by a factor of two with high-statistical data for the Dalitz distribution of $\eta \rightarrow 3\pi$ (both charged and neutral decays). More importantly, highly-boosted η production in JEF will offer improved systematics than the KLOE-II and A2-MAMI results that have much lower energies of η 's. In combination with a new Primakoff measurement of $\Gamma(\eta \rightarrow \gamma\gamma)$ from the on-going PrimEx-eta experiment (E-10-011) [24] to normalize the $\eta \rightarrow 3\pi$ decay width, it will allow an independent cross-check on the systematic uncertainty of the extracted quark mass ratio \mathcal{Q} for the first time.

Compared to the original JEF proposals [1, 2] submitted to the previous PACs, the scope of the JEF physics has been expanded mainly in two areas: (1) new physics searches have been broadened by not only searching for a leptophobic dark vector boson (B') [9] but also including dark photon [10, 11, 12, 13], hadrophilic scalar [14], and Axion-Like Particles (ALP) [25, 26, 27], probing two out of three dimension-four portals to the dark sector; (2) production of η' simultaneously with η at the similar rate will extend the mass coverage of new mediator search to ~ 1 GeV.

About 85% of matter in the Universe is Dark Matter (DM) whose constituents and interactions are unknown other than its gravitational properties. The stability of Dark Matter (DM) suggests that there may be a dark sector consisting of a rich symmetry structure with new forces and new particles. Dark sector may include one or more mediator particles coupled to the SM via a portal. The gauge and Lorentz symmetries of the Standard Model (SM) greatly restrict the ways in which the mediator can couple to the SM. There are three dimension-4 portals: the vector, scalar/pseudoscalar and fermion portals from the SM sector into the dark sector. Over the past decades, intensive efforts at the Large Hadron Collider (LHC) and underground laboratories have born no fruit for Weakly Interacting Massive Particles (WIMP), the simplest possible model for dark matter. There is a strong consensus among the physics community about the vital importance of broadening the scope of searches [28, 29, 30], both in the parameter space and in experimental approaches. The top-down models predict light mediators below GeV scale [31, 32]. These light states would have escaped detection thus far if they are very weakly coupled to the Standard Model. Recently, sub-GeV mediators have gained strong motivation, driven partly by several observed anomalies. These include several reported excesses in high-energy cosmic rays, which could be explained by dark matter annihilation [33, 34]. Scalar- or vector-mediated dark forces can also explain long-standing issues with galactic rotation curves and can solve small scale structure anomalies in dwarf galaxies and subhalos, while satisfying constraints on larger galaxy and cluster scales [35, 36, 37]. Lastly, there are the muon $g - 2$ anomaly [15, 10, 38] and an anomalous e^+e^- resonance observed in ^8Be decay [39, 40]. If these phenomena are interpreted in terms of new physics, all point toward mediator

there is another several a few words down 10

	Number: 1 Author: zisis	Subject: Cross-Out	Date: 2020-06-01, 8:18:21 AM	
	Number: 2 Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:20:23 AM	
	A probe of			
	Number: 3 Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:21:15 AM	
	their			
	Number: 4 Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:22:01 AM	
	-			
	Number: 5 Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:22:04 AM	
	s			
	Number: 6 Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:22:09 AM	
	-			
	Number: 7 Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:26:26 AM	
	The d			
	Number: 8 Author: zisis	Subject: Cross-Out	Date: 2020-06-01, 8:27:06 AM	
	Number: 9 Author: zisis	Subject: Replacement Text	Date: 2020-06-01, 8:28:02 AM	
	is			
	Number: 10	Author: zisis	Subject: Callout	Date: 2020-06-01, 8:27:32 AM
	there is another several a few words down			

References

- [1] L. Gan et al., the updated JEF proposal to JLab PAC45, 2017, https://www.jlab.org/exp_prog/proposals/17/C12-14-004.pdf.
- [2] L. Gan et al., the JEF proposal to JLab PAC42, 2014, https://www.jlab.org/exp_prog/proposals/14/PR12-14-004.pdf.
- [3] P. Adlarson [A2 Collaboration], EPJ Web Conf. **218**, 03002 (2019).
- [4] V. L. Kashevarov *et al.* [A2 Collaboration], Phys. Rev. Lett. **118**, no. 21, 212001 (2017) [arXiv:1701.04809 [nucl-ex]].
- [5] N. Hüsken *et al.* [WASA-at-COSY Collaboration], EPJ Web Conf. **199**, 01006 (2019).
- [6] S. S. Fang [BESIII Collaboration], PoS CD **15**, 032 (2016).
- [7] M. Berlowski [KLOE-2 Collaboration], EPJ Web Conf. **218**, 08003 (2019).
- [8] C. Gatto [REDTOP Collaboration], arXiv:1910.08505 [physics.ins-det].
- [9] S. Tulin, Phys.Rev. D89, 114008 (2014).
- [10] P. Fayet, Phys. Rev. D **75**, 115017 (2007) [hep-ph/0702176 [HEP-PH]].
- [11] M. Reece and L. T. Wang, JHEP **0907**, 051 (2009) [arXiv:0904.1743 [hep-ph]].
- [12] J. D. Bjorken, R. Essig, P. Schuster and N. Toro, Phys. Rev. D **80**, 075018 (2009) [arXiv:0906.0580 [hep-ph]].
- [13] B. Batell, M. Pospelov and A. Ritz, Phys. Rev. D **79**, 115008 (2009) [arXiv:0903.0363 [hep-ph]].
- [14] B. Batell, A. Freitas, A. Ismail and D. McKeen, Phys. Rev. D **100**, 095020 (2019) [arXiv:1812.05103 [hep-ph]].
- [15] Y. S. Liu, I. C. Cloët and G. A. Miller, Nucl. Phys. B , 114638 (2019).
- [16] B. A. Dobrescu, G. L. Landsberg and K. T. Matchev, Phys. Rev. D **63**, 075003 (2001) [arXiv:hep-ph/0005308 [hep-ph]].
- [17] D. Aloni, Y. Soreq and M. Williams, Phys. Rev. Lett. **123**, 031803 (2019) [arXiv:1811.03474 [hep-ph]].
- [18] M. Ramsey-Musolf, *et. al.*, phys. Rev., D63, 076007 (2001).
- [19] L. Ametller, J. Bijnens, and F. Cornet, Phys. Lett., B **276**, 185 (1992).
- [20] M. Knecht and A. Nyffeler, Phys. Rev. D **65**, 073034 (2002) doi:10.1103/PhysRevD.65.073034 [hep-ph/0111058].
- [21] G. Colangelo, S. Lanz, H. Leutwyler and E. Passemar, Eur. Phys. J. C **78**, no. 11, 947 (2018) [arXiv:1807.11937 [hep-ph]].
- [22] A. Anastasi *et al.* [KLOE-2 Collaboration], JHEP **1605**, 019 (2016) [arXiv:1601.06985 [hep-ex]].

This page contains no comments

- [23] S. Prakhov *et al.* [A2 Collaboration], Phys. Rev. C **97**, 065203 (2018) [arXiv:1803.02502 [hep-ex]].
- [24] A. Gasparian and L. Gan *et al.*, JLab proposal “A Precision Measurement of the η Radiative Decay Width via the Primakoff Effect”, https://www.jlab.org/exp_prog/proposals/10/PR12-10-011.pdf.
- [25] M. Bauer, M. Neubert and A. Thamm, JHEP **1712**, 044 (2017) [arXiv:1708.00443 [hep-ph]].
- [26] J. Hewett, *et al.* arXiv:1205.2671 [hep-ex].
- [27] W. Altmannshofer, S. Gori and D. J. Robinson, Phys. Rev. D **101**, 075002 (2020) [arXiv:1909.00005 [hep-ph]].
- [28] R. Essig *et al.*, arXiv:1311.0029 [hep-ph].
- [29] J. Alexander *et al.*, arXiv:1608.08632 [hep-ph].
- [30] M. Battaglieri *et al.*, arXiv:1707.04591 [hep-ph].
- [31] P. Fayet, Nucl. Phys. B **187**, 184 (1981).
- [32] M. Goodsell, J. Jaeckel, J. Redondo and A. Ringwald, JHEP **0911**, 027 (2009) [arXiv:0909.0515 [hep-ph]].
- [33] N. Arkani-Hamed, D. P. Finkbeiner, T. R. Slatyer and N. Weiner, Phys. Rev. D **79**, 015014 (2009) [arXiv:0810.0713 [hep-ph]].
- [34] M. Pospelov and A. Ritz, Phys. Lett. B **671**, 391 (2009) [arXiv:0810.1502 [hep-ph]].
- [35] S. Tulin and H. B. Yu, Phys. Rept. **730**, 1 (2018) [arXiv:1705.02358 [hep-ph]].
- [36] S. Tulin, H. Yu, K.M. Zurek, Phys. Rev. Lett. **110**, 111301 (2013).
- [37] S. Tulin, H. Yu, K.M. Zurek, Phys. Rev. D **87**, 115007-1 (2013).
- [38] M. Pospelov, Phys. Rev. D **80**, 095002 (2009) [arXiv:0811.1030 [hep-ph]].
- [39] A. J. Krasznahorkay *et al.*, Phys. Rev. Lett. **116**, 042501 (2016) [arXiv:1504.01527 [nucl-ex]].
- [40] J. L. Feng, B. Fornal, I. Galon, S. Gardner, J. Smolinsky, T. M. P. Tait and P. Tanedo, Phys. Rev. Lett. **117**, 071803 (2016) [arXiv:1604.07411 [hep-ph]].
- [41] M. Pospelov, “Light Scalar and Vector Extensions of the Standard Model”, Physics Seminars at JLab, May 16, 2014.

This page contains no comments