Data Analysis and Priliminary Results of the Proton Charge Radius (PRad) Experiment at JLab ¹

Weizhi Xiong, Chao Peng

Duke University

For the PRad Collaboration

In order to investigate the proton radius puzzle, the PRad experiment $(E12-11-106^2)$ was performed in 2016 in Hall B at Jefferson Lab, with 1.1 and 2.2 GeV unpolarized electron beam on a windowless H_2 gas flow target. The experiment aims to extract the electric form factor of the proton in an unprecedented low four-momentum transfer squared region, $Q^2 = 2 \times 10^{-4} - 0.06 \ (\text{GeV/c})^2$, with a sub-percent precision. The PRad experiment utilizes a non-magnetic calorimetric method with a high efficiency and high resolution calorimeter (HyCal), and two large area, high spatial resolution Gas Electron Multiplier (GEM) detectors. To control the systematic uncertainties, the absolute e - p elastic scattering cross sections are normalized to that of the well-known Møller scattering process, which is measured simultaneously within similar kinematics and experimental acceptances. Both the luminosity and the energy independent part of the detector acceptance and efficiency are cancelled in the ratio. In this talk, we will discuss details of the data analysis and present the physics results from the 2.2 GeV data.

¹This work is supported in part by the U.S. Department of Energy under Contract No. DE-FG02-03ER41231, NSF MRI award PHY-1229153, Thomas Jefferson National Accelerator Facility and Duke University

²Spokespersons: A. Gasparian (contact), H. Gao, M. Khandaker, D. Dutta