

Data Analysis and Preliminary Results of the Proton Charge Radius Experiment (PRad) at JLab ¹

Weizhi Xiong, Chao Peng

Duke University

For the PRad Collaboration

In order to investigate the proton radius puzzle, the PRad experiment (E12-11-106²) was performed in 2016 in Hall B at Jefferson Lab, with both 1.1 and 2.2 GeV unpolarized electron beams. The experiment aims to measure the $e - p$ elastic scattering cross section in an unprecedented low squared momentum transfer region ($Q^2 = 2 \times 10^{-4} - 0.06 \text{ (GeV/c)}^2$), with a sub-percent precision. The PRad experiment utilizes a non-magnetic calorimetric method with a large acceptance and high resolution calorimeter (HyCal), and two large area, high spatial resolution Gas Electron Multiplier (GEM) detectors. To have a better control over the systematic uncertainties, the absolute $e - p$ elastic scattering cross section is normalized to that of the well-known Møller scattering process, which is measured simultaneously within similar kinematics and geometrical acceptances; The windowless H₂-gas-flow target utilized in the experiment largely removes a typical background source, the target cell windows. In this talk, we will discuss details of the data analysis and present preliminary results from both beam energy settings.

¹This work is supported in part by the U.S. Department of Energy under Contract No. DE-FG02-03ER41231, NSF MRI award PHY-1229153, Thomas Jefferson National Accelerator Facility and Duke University

²Spokespersons: A. Gasparian (contact), H. Gao, M. Khandaker, D. Dutta