

A new direct detection electron scattering experiment to search for the X17 particle.

D. Dutta,¹ H. Gao,² A. Gasparian,³ T. J. Hague,^{3,4} N. Liyanage,⁵ R. Paremuzyan,⁶ C. Peng,⁷ W. Xiong,⁸ P. Achenbach,⁶ A. Ahmidouch,³ S. Ali,⁵ H. Avakian,⁶ C. Ayerbe-Gayoso,¹ X. Bai,⁵ M. Battaglieri,⁹ H. Bhatt,¹ A. Bianconi,^{10,11} J. Boyd,⁵ D. Byer,² P. L. Cole,¹² G. Costantini,^{10,11} S. Davis,³ M. De Napoli,¹³ R. De Vita,⁹ B. Devkota,¹ B. Dharmasena,⁵ J. Dunne,¹ L. El Fassi,¹ V. Gamage,⁵ L. Gan,¹⁴ K. Gnanvo,⁶ G. Gosta,^{10,11} D. Higinbotham,⁶ C. Howell,² S. Jeffas,⁵ S. Jian,⁵ A. Karki,¹ B. Karki,² V. Khachatryan,² M. Khandaker,¹⁵ V. Kubarovsky,⁶ I. Larin,¹⁶ M. Leali,^{10,11} V. Mascagna,^{17,11} G. Matousek,² S. Migliorati,^{10,11} R. Miskimen,¹⁶ P. Mohanmurthy,¹ H. Nguyen,⁵ E. Pasyuk,⁶ A. Rathnayake,⁵ J. Rittenhouse West,⁴ A. Shahinyan,¹⁸ A. Smith,² S. Stepanyan,⁶ E. van Nieuwenhuizen,² L. Venturelli,^{10,11} B. Yu,² Z. Zhao,² and J. Zhou^{2,4}

¹Mississippi State University

²Duke University

³North Carolina A&T State University

⁴Lawrence Berkeley National Laboratory

⁵University of Virginia

⁶Thomas Jefferson National Accelerator Facility

⁷Argonne National Laboratory

⁸Shandong University

⁹Istituto Nazionale di Fisica Nucleare, Sezione di Genova

¹⁰Dipartimento di Ingegneria dell'Informazione, Università di Brescia

¹¹Istituto Nazionale di Fisica Nucleare, sezione di Pavia

¹²Lamar University

¹³Istituto Nazionale di Fisica Nucleare, Sezione di Catania

¹⁴University of North Carolina, Wilmington

¹⁵Energy Systems

¹⁶University of Massachusetts

¹⁷DiSAT, Università dell'Insubria

¹⁸Yerevan Physics Institute

(Dated: January 2023)

A new electron scattering experiment (E12-21-003) [1] to verify and understand the nature of hidden sector particles, with particular emphasis on the so-called X17 particle, has been approved at Jefferson Lab. The search for these particles is motivated by new hidden sector models introduced to account for a variety of experimental and observational puzzles: excess in e^+e^- pairs observed in multiple nuclear transitions, the 4.2σ disagreement between experiments and the standard model prediction for the muon anomalous magnetic moment, and the small-scale structure puzzle in cosmological simulations. The aforementioned X17 particle has been hypothesized to account for the excess in e^+e^- pairs observed from the ^8Be M1, ^4He M0, and, most recently, ^{12}C E1 nuclear transitions to their ground states observed by the ATOMKI group. This experiment will use a high resolution electromagnetic calorimeter to search for or set new limits on the production rate of the X17 and other hidden sector particles in the 3 – 60 MeV mass range via their e^+e^- decay (or $\gamma\gamma$ decay with limited tracking). In these models, the 1 – 100 MeV mass range is particularly well-motivated and the lower part of this range still remains unexplored. Our proposed direct detection experiment will use a magnetic-spectrometer-free setup (the PRad apparatus) to detect all three final state particles in the visible decay of a hidden sector particle allowing for an effective control of the background and will cover the proposed mass range in a single setting. The use of the well-demonstrated PRad setup allows for an essentially ready-to-run and uniquely cost-effective search for hidden sector particles in the 3 – 60 MeV mass range with a sensitivity of 8.9×10^{-8} - 5.8×10^{-9} to ϵ^2 , the square of the kinetic mixing interaction constant between hidden and visible sectors.

I. INTRODUCTION

On large distance scales, the structure of the Universe as inferred from cosmological data is consistent with dark matter particles that are cold, collision-less, and interact with ordinary matter purely via gravity [2]. Decades of cosmological data have converged on the standard model of cosmology, dubbed ΛCDM , with cold dark matter (CDM) as a crucial ingredient [3] along with dark energy (the cosmological constant of general relativity, Λ)

and ordinary matter. Weakly interacting massive particles (WIMPs) have long been a primary candidate to explain dark matter (DM) [4, 5]. However, recent searches have ruled out a large segment of the phase space for WIMPs [6, 7] leading to new theoretical developments [8]. Additionally, several anomalous results in recent years have pointed to holes in our understanding of the nature of matter, discussed further in Sec. II.

These null results and the anomalous results have led to an increased interest in dark mediator models. In typi-

cal models these mediator particles (X) interact with the standard model (SM) through a ‘kinetic mixing’ mechanism creating a portal between the hidden sector and the SM. In most models the X particle couples to the electric charges of SM matter. Some newer models propose that the coupling to SM is tied to flavor of quarks or leptons and not via the electric charge. A common thread between many of the observed anomalies is that they can be explained by a new MeV-scale dark mediator particle. Given the convergence of such models on the 1 – 100 MeV mass range for a new particle, it is critical to focus experimental searches to this mass range in order to resolve the anomalies.

The availability of a high duty factor, high luminosity electron beam at Jefferson Lab provides an ideal setup to search for MeV-scale dark mediators with small coupling constants. The well tested PRad setup in Hall B [9] will be used in this experiment to reach our physics goals. Using a magnetic-spectrometer-free setup allows the experiment to be sensitive to the full mass range in a single experimental setting. The detection of all three final state particles in the PbWO_4 calorimeter along with tracking with GEM chambers allows for an effective control of the backgrounds. Moreover, it provides an essentially ready-to-run and uniquely cost effective search for hidden-sector particles in the 3 - 60 MeV mass range.

II. PHYSICS MOTIVATION

The primary motivation for this new experiment is to resolve the X17 anomaly. A 2015 experiment [10] at the ATOMKI 5 MV Van de Graaff accelerator sought to study a previous 9 MeV anomaly [11–13]. While this experiment nullified and explained the previous result, it saw an excess of e^+e^- pairs beyond the expectation of internal pair creation (IPC) for a different mass. A subsequent analysis of these results has shown that the 6.8σ anomaly is consistent with a new particle with a mass of 16.84 MeV, dubbed X17. A followup experiment by the ATOMKI group was conducted on the 20.01 MeV $0^- \rightarrow 0^+$ transition in ^4He . The results for ^4He also reports an e^+e^- excess consistent with the so-called X17 particle [14, 15]. In a new 2022 experiment, the group has reported the anomaly in the ^{12}C E1 transition [16]. Of note, this most recent result agrees with the previous masses but disagrees with the measured branching ratios.

Feng *et al.* [18] analyzed the X17 signal against existing constraints. The proposed explanation is that the signal is from the decay of a protophobic gauge boson that mediates a fifth force with a length scale of 12 fm. This explanation can also possibly explain the muon anomalous magnetic moment and an excess of $\pi^0 \rightarrow e^+e^-$ decays [19]. The current constraints from leptonic production mechanisms, where the effective coupling to a new force-carrier is proportional to electric charge, are shown in Fig. 1. For more generic fifth-force models with quark flavor-dependent couplings [18], a much wider parameter




FIG. 1. Current constraints on a fifth force explanation of the ^8Be anomaly. The vertical axis is the leptonic coupling strength relative to α_{QED} , with horizontal axis the mass of the mediator. Excluded regions, in gray, are taken from measurements that depend solely on leptonic interactions. Dark photon exclusions via hadronic measurements are not shown. Reproduced from Ref. [17].

space with multiple couplings must be considered.

The fifth force based explanations are being challenged by recent reanalyses [20, 21] and the observed discrepancies could be the result of as-yet-unidentified nuclear reactions, excited diquark states, or experimental effects. Nonetheless, these results have garnered a lot of attention, and must be independently validated with the highest urgency.

The Muon $g - 2$ collaboration has recently reported their measurement of the muon magnetic moment at Fermi National Accelerator Laboratory (FNAL) which is consistent with their previous measurement at Brookhaven National Laboratory (BNL) [22, 23]. These two results, with their uncertainties combined, show a 4.2σ deviation from the Standard Model prediction.

There have been several proposals as to new physics that could lead to this deviation. One such class of explanations shows that undiscovered hidden sector particles can directly contribute to loop corrections in the calculation of $(g - 2)_\mu$. The contributions of a hidden sector particle to the $(g - 2)_\mu$ scales as $1/m_X^2$, thus a smaller mass will have a comparably larger contribution than a higher mass particle. There have been numerous publications proposing new particles (such as dark photons) in the MeV-range that, at a minimum, partially account for the muon anomalous magnetic moment [24–27]. Fig. 1 shows that there is an overlap in the phase space for a particle that would simultaneously resolve the X17 and $(g - 2)_\mu$ anomalies.

Several recent models have suggested that the coupling of the X17 to SM is tied to flavor of quarks or leptons and not via the electric charge. One such example is the protophobic X17 proposed in Ref. [18] used to explain the X17 anomaly. If an X17 were to have flavor-dependent couplings the parameter space for it is more open than

previously thought [28]. These new possibilities point to an urgent need for independent verification of the X17 anomaly.

III. EXPERIMENTAL METHOD AND SETUP

This experiment will focus on the bremsstrahlung-like production of hidden sector particles from the initial electron or the scattered electron (both shown in Fig.2), in the 3–60 MeV mass range. When searching for new particles, it is of the utmost importance to minimize backgrounds in order to prevent false “bumps” in the mass spectrum. The primary QED background in this experiment (see Fig. 3) are from the radiative pair production which is an irreducible background and the Bethe-Heitler trident reactions which can be kinematically suppressed.




FIG. 2. Bremsstrahlung-like production of a hidden sector force carrier X from electron scattering. The left diagram shows production from the incoming electron and the right diagram shows production from the scattered electron.




FIG. 3. The QED background from radiative (left and middle) and Bethe-Heitler (right) process.

The experiment plans to reuse the PRad setup (shown in Fig. 4), with the addition of a Tantalum foil target placed 7.5 m upstream of the calorimeter. The high resolution PbWO_4 crystal part of the HyCal electromagnetic calorimeter [29, 30] will be used together with a new flash ADC (FADC) based readout system for the calorimeter. A 2.2 and 3.3 GeV Continuous Wave (CW) electron beam from CEBAF will be incident on a retractable ultra-thin target consisting of a $1\ \mu\text{m}$ Tantalum foil. The scattered particles will traverse the 7.5 m long flight path in vacuum, en route to a pair of common ionization volume GEM chambers coupled to the HyCal EM calorimeter. The vacuum chamber will consist of the PRad target chamber (or an appropriate diameter beam

pipe) and the 5 m long PRad vacuum chamber with a thin window. The vacuum chamber matches the geometrical acceptance of the calorimeter. An extension piece will be added to the upstream end to couple the PRad target chamber to the super-harp which will now hold the target foil. A reducer ring will be attached to the downstream exit of the PRad vacuum chamber and a new 1 mm thick Al exit window will be used such that it matches the PbWO_4 portion of the calorimeter. Two layers of GEM detectors will add a modest tracking capability to help reduce the photon background and to reduce the background originating from the vacuum chamber exit window.




FIG. 4. Schematic of the experimental setup.

The experiment will use a “bump hunt” technique in the direct detection search of heretofore unknown MeV mass particles. All 3 cluster events with individual cluster energy within $(0.02-0.85) \times E_{beam}$ and with the sum of total energy deposited $E_{sum} > 0.7 \times E_{beam}$ will be recorded and examined for “bumps” in the M_{e+e-} (or $M_{\gamma\gamma}$) invariant mass spectrum reconstructed from these events. This will allow for the X -particles production by virtual photons over a wide energy range in the forward solid angle coverage of the PbWO_4 calorimeter. The capability of detecting events produced in a wide energy and angle range, in a single experimental setting, is one of the important features of our experiment which stands in contrast to all other magnetic spectrometer methods. The tracking provided by the pair of GEM chambers will be used to suppress background events from the experimental setup, particularly, the large area window at the exit of the vacuum chamber. The GEM chambers will also be used to veto any neutral particles. Only the PbWO_4 part of the HyCal calorimeter will be used in the experiment to ensure high invariant mass resolution. The experimental method discussed here applies directly to any spin-1 boson in the hidden sector that decay directly to a lepton pair.

IV. PROJECTED RESULTS

In order to estimate the projected sensitivity to ϵ^2 , a comprehensive simulation of the experiment was carried out using the Geant4 simulation package developed for the PRad experiment [9]. The simulation used a hybrid




FIG. 5. Acceptance of the spectrometer setup for both 2.2 and 3.3 GeV beam energies.




FIG. 6. Invariant mass resolution of the spectrometer for 2.2 GeV beam energy.

event generator that was built by combining the Madgraph5 generator [34] for the Bethe-Heitler process with the Geant4 generator (with Bethe-Heitler turned off) to utilize the best features of the two generators. The background was simulated for 200 s of beamtime and then scaled by sampling the number of events bin-by-bin to

give the expected background for 30 days of 100 nA, 3.3 GeV beam.

The production rates for the X particle are calculated using the rate equation (Eq. 14) from Ref. [35] and rearranged to give the sensitivity as:

$$\epsilon^2 = \frac{N_X}{5 \times N_e T \frac{m_e^2}{m_X^2}}, \quad (1)$$

where N_X is the number of X particles produced, N_e is the number of incident electrons (1.62×10^{18} for 30 days at 100 nA), T is the target thickness (2.5×10^{-4} r.l.), m_X is the mass of the produced X , and ϵ^2 is the square of the dimensionless coupling constant of the X to SM matter. The lowest ϵ^2 achievable is calculated using Eq. 1 with N_X obtained by scaling the signal counts with the detection efficiency and impact of the decay length (shown in Fig. 5). The background event count is calculated by selecting events from the background simulation within a $\pm 3\sigma_{m_X}$ window. The typical invariant mass resolution σ_{m_X} obtained from the simulation is shown in Fig. 6. The signal counts necessary given the background counts are calculated using the criteria for a 2.3σ significance used by other experiments (Eq. 2).

$$\frac{N_{\text{signal}}}{\sqrt{N_{\text{signal}} + N_{\text{bgd}}}} \geq 2.3. \quad (2)$$




FIG. 7. Projected coverage of the square of the coupling constant (ϵ^2) and mass (m_X) parameter space by this proposed experiment is shown as the thick red lines for the combined statistics of the two beam energies. The projections are superimposed on top of a constraints plot adapted from Ref. [8], with additional data from Refs.[31–33].

The 2.2 GeV runs will serve as reference runs to better understand the background and the signal, so it was conservatively estimated that 50% of their statistics can be combined into the 3.3 GeV data for the final results. Using these range of sensitivities the bounds for the $\epsilon^2 - m_X$ parameter space is plotted in Fig. 7 for the combined projected statistics.

V. CONCLUSION

The X17 anomaly suggests the existence of a new hidden sector particle. It is of critical importance to resolve this anomaly quickly, as it not only could point to new physics but also has the potential to resolve additional anomalies. In this white paper, we have described a nearly ready-to-run experiment at JLab using the PRad spectrometer that has the capability to fully address this anomaly. By using a magnet-free spectrometer, two beam energies, and detecting all three final state the experiment will provide unprecedented control of the experimental background, leading to a results with uniquely low systematics.

-
- [1] A. Ahmidouch *et al.*, A Direct Detection Search for Hidden Sector New Particles in the 3-60 MeV Mass Range, (2021), arXiv:2108.13276 [nucl-ex].
 - [2] T. S. and Y. H.-B., Dark Matter Self-interactions and Small Scale Structure, *Phys. Rep.* **730**, 1 (2018), arXiv:1705.02358 [hep-ph].
 - [3] B. N. A., O. J. P., P. S., and S. P. J., The Cosmic Triangle: Revealing the State of the Universe, *Science* **284**, 1481 (1999), arXiv:9906463 [astro-ph].
 - [4] G. Jungman, M. Kamionkowski, and K. Griest, Supersymmetric dark matter, *Physics Reports* **267**, 195 (1996).
 - [5] J. L. Feng, Dark Matter Candidates from Particle Physics and Methods of Detection, *Ann. Rev. Astron. Astrophys.* **48**, 495 (2010), arXiv:1003.0904 [astro-ph.CO].
 - [6] M. Battaglieri *et al.*, US Cosmic Visions: New Ideas in Dark Matter 2017: Community Report, *U.S. Cosmic Visions: New Ideas in Dark Matter*, arXiv:1707.04591 (2017), arXiv:1707.04591 [hep-ph].
 - [7] E. Aprile *et al.*, Dark Matter Search Results from a One Ton-Year Exposure of XENON1T, *Phys. Rev. Lett.* **121**, 111302 (2018).
 - [8] J. Alexander *et al.*, Dark Sectors 2016 Workshop: Community Report, (2016), arXiv:1608.08632 [hep-ph].
 - [9] W. Xiong *et al.*, A small proton charge radius from an electron-proton scattering experiment, *Nature* **575**, 147 (2019).
 - [10] A. J. Krasznahorkay *et al.*, Observation of Anomalous Internal Pair Creation in Be8 : A Possible Indication of a Light, Neutral Boson, *Phys. Rev. Lett.* **116**, 042501 (2016), arXiv:1504.01527 [nucl-ex].
 - [11] F. de Boer, O. Fröhlich, K. Stiebing, K. Bethge, H. Bokemeyer, A. Balanda, A. Buda, R. van Dantzig, T. Elze, H. Folger, J. van Klinken, K. Müller, K. Stelzer, P. Thee, and M. Waldschmidt, A deviation in internal pair conversion, *Phys. Lett. B* **388**, 235 (1996).
 - [12] F. W. N. de Boer, R. van Dantzig, J. van Klinken, K. Bethge, H. Bokemeyer, A. Buda, K. A. Müller, and K. E. Stiebing, Excess in nuclear e^+e^- pair near 9 MeV/c² invariant mass, *J. Phys. G* **23**, L85 (1997).
 - [13] F. W. N. de Boer, K. Bethge, H. Bokemeyer, R. van Dantzig, J. van Klinken, V. Mironov, K. A. Muller, and K. E. Stiebing, Further search for a neutral boson with a mass around 9-MeV/c², *J. Phys. G* **27**, L29 (2001), arXiv:hep-ph/0101298.
 - [14] A. J. Krasznahorkay *et al.*, New evidence supporting the existence of the hypothetical X17 particle, (2019), arXiv:1910.10459 [nucl-ex].
 - [15] A. J. Krasznahorkay, M. Csatlós, L. Csige, J. Gulyás, A. Krasznahorkay, B. M. Nyakó, I. Rajta, J. Timár, I. Vajda, and N. J. Sas, New anomaly observed in He4 supports the existence of the hypothetical X17 particle, *Phys. Rev. C* **104**, 044003 (2021), arXiv:2104.10075 [nucl-ex].
 - [16] A. J. Krasznahorkay *et al.*, New anomaly observed in ¹²C supports the existence and the vector character of the hypothetical X17 boson, *Phys. Rev. C* **106**, L061601 (2022), arXiv:2209.10795 [nucl-ex].
 - [17] Search for new physics in e+e-final states near an invariant mass of 17 mev using the cebaf injector, http://www.jlab.org/exp_prog/proposals/20/PR12-20-001_Proposal.pdf (2020).
 - [18] J. L. Feng, B. Fornal, I. Galon, S. Gardner, T. M. P. Smolinsky, J. and Tait, and P. Tanedo, Protophobic Fifth-Force Interpretation of the Observed Anomaly in ⁸Be Nuclear Transitions, *Phys. Rev. Lett.* **117**, 071803 (2016), arXiv:1604.07411 [hep-ph].
 - [19] E. Abouzaid *et al.* (KTeV), Measurement of the Rare Decay $\pi^0 \rightarrow e^+e^-$, *Phys. Rev. D* **75**, 012004 (2007), arXiv:hep-ex/0610072.
 - [20] X. Zhang and G. A. Miller, Can a protophobic vector boson explain the ATOMKI anomaly?, *Phys. Lett. B* **813**, 136061 (2021), arXiv:2008.11288 [hep-ph].
 - [21] V. Kubarovsky, J. R. West, and S. J. Brodsky, Quantum Chromodynamics Resolution of the ATOMKI Anomaly in ⁴He Nuclear Transitions, (2022), arXiv:2206.14441 [hep-ph].
 - [22] T. Albahri *et al.* (Muon g-2), Magnetic-field measurement and analysis for the Muon $g - 2$ Experiment at Fermilab, *Phys. Rev. A* **103**, 042208 (2021), arXiv:2104.03201 [hep-ex].
 - [23] G. W. Bennett *et al.* (Muon g-2), Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL, *Phys. Rev. D* **73**, 072003 (2006), arXiv:hep-ex/0602035.
 - [24] G. Arcadi, A. S. De Jesus, T. B. De Melo, F. S. Queiroz, and Y. S. Villamizar, A 2HDM for the g-2 and Dark Matter, arXiv:2104.04456 (2021), arXiv:2104.04456 [hep-ph].

- [25] P. Athron, C. Balázs, D. H. Jacob, D. Kotlarski, W. and Stöckinger, and H. Stöckinger-Kim, New physics explanations of a_μ in light of the FNAL muon $g - 2$ measurement, arXiv:2104.03691 (2021), arXiv:2104.03691 [hep-ph].
- [26] D. Borah, M. Dutta, S. Mahapatra, and N. Sahu, Muon ($g - 2$) and XENON1T Excess with Boosted Dark Matter in $L_\mu - L_\tau$ Model, arXiv:2104.05656 (2021), arXiv:2104.05656 [hep-ph].
- [27] S.-F. Ge, X.-D. Ma, and P. Pasquini, Probing the Dark Axion Portal with Muon Anomalous Magnetic Moment, arXiv:2104.03276 (2021), arXiv:2104.03276 [hep-ph].
- [28] J. L. Feng *et al.*, The Forward Physics Facility at the High-Luminosity LHC, (2022), arXiv:2203.05090 [hep-ex].
- [29] A. Gasparian (PrimEx), Performance of PWO crystal detectors for a high resolution hybrid electromagnetic calorimeter at Jefferson Lab, in *10th International Conference on Calorimetry in High Energy Physics (CALOR 2002)* (2002) pp. 208–214.
- [30] A. Gasparian (PrimEx), A high performance hybrid electromagnetic calorimeter at Jefferson Lab, in *11th International Conference on Calorimetry in High-Energy Physics (Calor 2004)* (2004) pp. 109–115.
- [31] R. Aaij *et al.* (LHCb), Search for Dark Photons Produced in 13 TeV pp Collisions, Phys. Rev. Lett. **120**, 061801 (2018), arXiv:1710.02867 [hep-ex].
- [32] R. Aaij *et al.* (LHCb), Search for $A' \rightarrow \mu^+ \mu^-$ Decays, Phys. Rev. Lett. **124**, 041801 (2020), arXiv:1910.06926 [hep-ex].
- [33] N. Baltzell *et al.*, The Heavy Photon Search Experiment, (2022), arXiv:2203.08324 [hep-ex].
- [34] E. Conte, B. Fuks, and G. Serret, MadAnalysis 5, A User-Friendly Framework for Collider Phenomenology, Comput. Phys. Commun. **184**, 222 (2013), arXiv:1206.1599 [hep-ph].
- [35] J. D. Bjorken, R. Essig, P. Schuster, and N. Toro, New Fixed-Target Experiments to Search for Dark Gauge Forces, Phys. Rev. D **80**, 075018 (2009), arXiv:0906.0580 [hep-ph].